

A class of relations among multiple zeta values

Gaku Kawashima

Graduate School of Mathematics,

Nagoya University, Chikusa-ku, Nagoya464-8602, Japan

E-mail: m02009c@math.nagoya-u.ac.jp

Abstract

We prove a new class of relations among multiple zeta values (MZV's) which contains Ohno's relation. We also give the formula for the maximal number of independent MZV's of fixed weight, under our new relations. To derive our formula for MZV's, we consider the Newton series whose values at non-negative integers are finite multiple harmonic sums.

keywords multiple zeta value, the Newton series, Ohno's relation

1 Introduction

Let $k_1, \dots, k_p$ be positive integers and let $k_p \geq 2$. Multiple zeta values (MZV's) are defined by

$$\zeta(k_1, \dots, k_p) = \sum_{0 < n_1 < \dots < n_p} \frac{1}{n_1^{k_1} \dots n_p^{k_p}}.$$

The sum $k_1 + \dots + k_p$ is called the weight of the multiple zeta value $\zeta(k_1, \dots, k_p)$. These numbers were considered first by Euler [2]; he studied the case $p = 2$. The general case was introduced in [4] and [19]. In recent years, many researchers have studied these numbers in connection with Galois representations, arithmetic geometry, quantum groups, invariants for knots, mathematical physics, etc. Many $\mathbb{Q}$ -linear relations are known among these values, e.g. Ohno's relation, the cyclic sum formula and the derivation relation [7, 8, 15, 16]. The famous relation due to Ohno generalizes the duality, the sum formula and Hoffman's relation simultaneously. Ihara, Kaneko and Zagier investigated the regularized double shuffle relations in [8], where they conjectured that these relations imply all $\mathbb{Q}$ -linear relations. Recently in [10] Kaneko proposed a conjectural generalization of the derivation relation for MZV's and in [17] Tanaka proved this conjecture by reducing it to the relations studied in this paper. The aim of this paper is to give a new class of relations among MZV's. We shall prove that our new class of relations contains Ohno's relation. We conjecture that it also contains the cyclic sum formula. We have checked this up to weight 12.

We shall outline how our relations among MZV's can be derived. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$ be a multi-index (i.e., an ordered set of positive integers) and n a

non-negative integer. We consider the finite multiple harmonic sums

$$s_{\boldsymbol{\mu}}(n) = \sum_{0 \leq n_1 \leq \dots \leq n_p = n} \frac{1}{(n_1 + 1)^{\mu_1} \dots (n_p + 1)^{\mu_p}}$$

and

$$\begin{aligned} S_{\boldsymbol{\mu}}(n) &= \sum_{0 \leq k < n} s_{\boldsymbol{\mu}}(k) \\ &= \sum_{0 \leq n_1 \leq \dots \leq n_p < n} \frac{1}{(n_1 + 1)^{\mu_1} \dots (n_p + 1)^{\mu_p}}. \end{aligned}$$

In the following, we denote the set of non-negative integers by $\mathbb{N} = \{0, 1, 2, \dots\}$. We define the operator ∇ on the space $\mathbb{C}^{\mathbb{N}}$ of complex-valued sequences by

$$(\nabla a)(n) = \sum_{k=0}^n (-1)^k \binom{n}{k} a(k)$$

for $a \in \mathbb{C}^{\mathbb{N}}$ and $n \in \mathbb{N}$. In Section 3, we shall prove our key formula for the finite multiple harmonic sums $s_{\boldsymbol{\mu}}(n)$ (Theorem 3.8). As a corollary of Theorem 3.8, we shall prove that

$$\nabla s_{\boldsymbol{\mu}} = s_{\boldsymbol{\mu}^*},$$

where $\boldsymbol{\mu}^*$ is some multi-index determined by $\boldsymbol{\mu}$. We note that this assertion was proved independently by Hoffman in [6, Theorem 4.2].

In Section 5, in order to derive our relations among MZV's, we consider the Newton series which interpolates the finite multiple harmonic sums $S_{\boldsymbol{\mu}}(n)$. This Newton series $F_{\boldsymbol{\mu}}(z)$ is defined by

$$F_{\boldsymbol{\mu}}(z) = \sum_{n=0}^{\infty} (-1)^n (\nabla S_{\boldsymbol{\mu}})(n) \binom{z}{n}, \quad \binom{z}{n} = \frac{z(z-1) \dots (z-n+1)}{n!},$$

where z is a complex number with $\operatorname{Re} z > -1$. In fact, for any $n \in \mathbb{N}$, we have $F_{\boldsymbol{\mu}}(n) = S_{\boldsymbol{\mu}}(n)$. It is well known that $S_{\boldsymbol{\mu}}(n)S_{\boldsymbol{\nu}}(n)$ is a $\mathbb{Z}$ -linear combination of $S_{\boldsymbol{\lambda}}(n)$'s:

$$S_{\boldsymbol{\mu}}(n)S_{\boldsymbol{\nu}}(n) = \sum_i c_i S_{\boldsymbol{\lambda}^i}(n), \quad c_i \in \mathbb{Z}.$$

The multi-indices $\boldsymbol{\lambda}^i$ and the coefficients c_i depend only on $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$, but not on n . From this fact, we derive the functional equations

$$F_{\boldsymbol{\mu}}(z)F_{\boldsymbol{\nu}}(z) = \sum_i c_i F_{\boldsymbol{\lambda}^i}(z), \tag{1}$$

which are valid for $\operatorname{Re} z > -1$. Since the derivatives $F_{\boldsymbol{\mu}}^{(m)}(0)$ ($m \geq 1$) can be expressed in terms of MZV's, quadratic relations among MZV's are derived from (1) (Corollary 5.4).

We mainly consider the first derivatives only. Since $F_{\boldsymbol{\mu}}(0)$ vanishes, equation (1) implies a linear relation

$$0 = \sum_i c_i F'_{\boldsymbol{\lambda}^i}(0).$$

In this way, we get a set of linear relations among MZV's which can be written explicitly (Corollary 5.5). We shall explain this formula. Let V_k be the $\mathbb{Q}$ -vector space whose basis is the set of all multi-indices of weight k and let $V = \bigoplus_{k \geq 1} V_k$. We define $\mathbb{Q}$ -linear mappings σ, u from V to V by

$$\begin{aligned} \sigma(\mu_1, \dots, \mu_p) &= (-1)^p (\mu_1, \dots, \mu_p), \\ u(\mu_1, \dots, \mu_p) &= \sum_{(\nu_1, \dots, \nu_q) \geq (\mu_1, \dots, \mu_p)} (\nu_1, \dots, \nu_q), \end{aligned}$$

where $(\nu_1, \dots, \nu_q)$ runs over all refinements of $(\mu_1, \dots, \mu_p)$ (see §2 for the precise definition). We define the $\mathbb{Q}$ -linear mapping ζ^+ from V to $\mathbb{R}$ by

$$\zeta^+(\mu_1, \dots, \mu_p) = \zeta(\mu_1, \dots, \mu_{p-1}, \mu_p + 1).$$

Then Corollary 5.5 asserts that

$$\zeta^+(u\sigma(\boldsymbol{\mu} * \boldsymbol{\nu})) = 0$$

for any multi-indices $\boldsymbol{\mu}$ and $\boldsymbol{\nu}$. (The multiplication $*$ on V is the harmonic product.) For example, for $\boldsymbol{\mu} = \boldsymbol{\nu} = (1)$, we have

$$\begin{aligned} \boldsymbol{\mu} * \boldsymbol{\nu} &= (2) + 2(1, 1), \\ \sigma(\boldsymbol{\mu} * \boldsymbol{\nu}) &= -(2) + 2(1, 1) \end{aligned}$$

and

$$u\sigma(\boldsymbol{\mu} * \boldsymbol{\nu}) = -\{(2) + (1, 1)\} + 2(1, 1) = -(2) + (1, 1).$$

Therefore we obtain

$$\zeta^+(2) = \zeta^+(1, 1) \quad (\text{i.e., } \zeta(3) = \zeta(1, 2)),$$

which is due to Euler.

Under the harmonic product $*$, the $\mathbb{Q}$ -vector space $\mathbb{Q} \oplus V$ becomes a commutative $\mathbb{Q}$ -algebra. This is isomorphic to the commutative $\mathbb{Q}$ -algebra of quasi-symmetric functions, which is known to be the polynomial algebra on the set of Lyndon words [12, Section 2], and whose enumeration is well known. From this fact, it is easily seen that

$$\begin{aligned} d(k) &:= \dim_{\mathbb{Q}}(u\sigma(V * V) \cap V_k) \\ &= 2^{k-1} - \frac{1}{k} \sum_{d|k} \mu\left(\frac{k}{d}\right) 2^d \quad (k \geq 2), \end{aligned} \tag{2}$$

where μ is the Möbius function. The right-hand side of (2) appeared in the recent research of Kaneko for the extended derivation relation, and this equation was suggested to the author by Kaneko.

We shall compare the dimensions of three subspaces of $\ker \zeta^+ \cap V_k$. We define the sequence $\{z_k\}_{k=1}^\infty$ by

$$z_1 = z_2 = z_3 = 1, \quad z_k = z_{k-2} + z_{k-3} \quad (k \geq 4)$$

and put $d_Z(k) = 2^{k-1} - z_k$. Then the dimension of the space $\ker \zeta^+ \cap V_k$ is conjectured to be equal to $d_Z(k)$ by Zagier [19]. We denote the dimension of the space $V_{Ohno} \cap V_k$ by $d_O(k)$, where V_{Ohno} is the subspace of $\ker \zeta^+$ which corresponds to Ohno's relation. The following table gives us the numerical values of $d_Z(k)$, $d(k)$ and $d_O(k)$ for $2 \leq k \leq 11$.

k	2	3	4	5	6	7	8	9	10	11
$d_Z(k)$	1	3	6	14	29	60	123	249	503	1012
$d(k)$	1	2	5	10	23	46	98	200	413	838
$d_O(k)$	1	2	5	10	23	46	98	199	411	830

The product of two MZV's is expressed as a $\mathbb{Z}$ -linear combination of MZV's in two ways; the harmonic product relations and the shuffle product relations. We conjecture that under the shuffle product relations, Corollary 5.4 for $m = 1, 2$ gives all $\mathbb{Q}$ -linear relations among MZV's. This conjecture is due to Tanaka. He checked that the dimension of the subspace of $\ker \zeta^+$ which corresponds to these linear relations is equal to $d_Z(k)$ for $2 \leq k \leq 11$.

2 Multi-indices

A finite sequence of positive integers is called a multi-index and we denote the set of all multi-indices by I . We denote by V the $\mathbb{Q}$ -vector space whose basis is I . For $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$, we call $l(\boldsymbol{\mu}) := p$ and $|\boldsymbol{\mu}| := \sum_{i=1}^p \mu_i$ the length and the weight of $\boldsymbol{\mu}$, respectively. For a positive integer m , we denote the set of all multi-indices of weight m by I_m . We define the mapping $\mathcal{S}_m$ for a positive integer m by

$$\mathcal{S}_m: I_m \rightarrow 2^{\{1,2,\dots,m-1\}}, \quad (\mu_1, \dots, \mu_p) \mapsto \left\{ \sum_{i=1}^k \mu_i \mid 1 \leq k < p \right\}.$$


This is a bijection. For example, we obtain

$$\mathcal{S}_5(2, 2, 1) = \{2, 4\} \quad \text{and} \quad \mathcal{S}_5(1, 1, 3) = \{1, 2\}$$

by the following diagrams



We define an order on I by setting $\mu \geq \nu$ if $|\mu| = |\nu|$ and $\mathcal{S}_\mu(\mu) \supset \mathcal{S}_\nu(\nu)$. In other words, we write $\mu \geq \nu$ if μ is a refinement of ν . For example, the order on I_4 is given by the following Hasse diagram



For $\mu = (\mu_1, \dots, \mu_p) \in I$ with $|\mu| = m$, we put

$$\begin{aligned}
 (\mu) &= \mu^ = \mathcal{S}_m^{-1}(\mathcal{S}_m(\mu)^c), \\
 \tau(\mu) &= (\mu_p, \dots, \mu_1), \\
 \sigma(\mu) &= (-1)^{l(\mu)} \mu, \\
 u(\mu) &= \sum_{\nu \geq \mu} \nu
 \end{aligned}$$


and

$$d(\mu) = \sum_{\nu \leq \mu} \nu.$$

($\mathcal{S}_m(\mu)^c$ denotes the complement of $\mathcal{S}_m(\mu)$ in $\{1, 2, \dots, m-1\}$.) We extend them as $\mathbb{Q}$ -linear mappings from V to V . All of these are bijections. We give examples of μ^* . We have

$$(1, 2, 3)^* = (2, 2, 1, 1), \quad (2, 2, 2)^* = (1, 2, 2, 1) \quad \text{and} \quad (4, 1, 1)^* = (1, 1, 1, 3)$$

by the diagrams



where the lower arrows are in the complementary slots to the upper arrows. We can also calculate these correspondences by the diagrams

$$\begin{array}{c}
 2 \ 2 \ 1 \ 1 \\
 1 \circ \\
 2 \circ \circ \\
 3 \ \circ \circ \circ
 \end{array}, \quad
 \begin{array}{c}
 1 \ 2 \ 2 \ 1 \\
 2 \circ \circ \\
 2 \ \circ \circ \\
 2 \ \circ \circ
 \end{array}
 \quad \text{and} \quad
 \begin{array}{c}
 1 \ 1 \ 1 \ 3 \\
 4 \circ \circ \circ \circ \\
 1 \ \circ \\
 1 \ \circ
 \end{array}.$$

It is easily seen that

$$l(\boldsymbol{\mu}) + l(\boldsymbol{\mu}^*) = |\boldsymbol{\mu}| + 1 \quad (3)$$

for any $\boldsymbol{\mu} \in I$. Let

$$\boldsymbol{\mu} = (\mu_1, \dots, \mu_r, \underbrace{1, \dots, 1}_l) \quad (\mu_r \geq 2, r \geq 0, l \geq 0)$$

and $\boldsymbol{\mu}^* = (\mu_1^*, \dots, \mu_q^*)$. Then we note that

$$\mu_q^* = l + 1. \quad (4)$$

Proposition 2.1. *We have $*\tau = \tau*$, $\sigma\tau = \tau\sigma$, $u\tau = \tau u$ and $d\tau = \tau d$.*

Proof. These are easily seen. $\square$

Lemma 2.2. *Let $T_1 \subset T_2$ be finite sets. Then we have*

$$\sum_{T_1 \subset S \subset T_2} (-1)^{\#S} = \begin{cases} (-1)^{\#T_2} & \text{if } T_1 = T_2 \\ 0 & \text{if } T_1 \subsetneq T_2. \end{cases}$$

Proof. The left-hand side equals

$$\begin{aligned} (-1)^{\#T_1} \sum_{S \subset (T_2 \setminus T_1)} (-1)^{\#S} &= (-1)^{\#T_1} \sum_{k=0}^{\#(T_2 \setminus T_1)} (-1)^k \binom{\#(T_2 \setminus T_1)}{k} \\ &= (-1)^{\#T_1} (1 - 1)^{\#(T_2 \setminus T_1)}. \end{aligned}$$

Thus we proved the lemma. $\square$

Proposition 2.3. *We have*

- (i) $*d* = u$,
- (ii) $d\sigma d\sigma = \text{id}_V$,
- (iii) $u\sigma u\sigma = \text{id}_V$,
- (iv) $d*d^{-1} = -u\sigma$,
- (v) $u^{-1}*u = -\sigma d$.

Proof. Let $\boldsymbol{\mu}$ be any element of I and let $|\boldsymbol{\mu}| = m$.

(i) We have

$$*d*(\boldsymbol{\mu}) = \sum_{\boldsymbol{\nu} \leq \boldsymbol{\mu}^*} \boldsymbol{\nu}^* = \sum_{\boldsymbol{\nu}^* \geq \boldsymbol{\mu}} \boldsymbol{\nu}^* = u(\boldsymbol{\mu}).$$

(ii) We have

$$d\sigma d\sigma(\boldsymbol{\mu}) = (-1)^{l(\boldsymbol{\mu})} \sum_{\boldsymbol{\nu} \leq \boldsymbol{\mu}} (-1)^{l(\boldsymbol{\nu})} \sum_{\boldsymbol{\lambda} \leq \boldsymbol{\nu}} \boldsymbol{\lambda} = (-1)^{l(\boldsymbol{\mu})} \sum_{\boldsymbol{\lambda} \leq \boldsymbol{\mu}} \left\{ \sum_{\boldsymbol{\lambda} \leq \boldsymbol{\nu} \leq \boldsymbol{\mu}} (-1)^{l(\boldsymbol{\nu})} \right\} \boldsymbol{\lambda}.$$

By Lemma 2.2, we have

$$\sum_{\boldsymbol{\lambda} \leq \boldsymbol{\nu} \leq \boldsymbol{\mu}} (-1)^{l(\boldsymbol{\nu})} = \sum_{\mathcal{S}_m(\boldsymbol{\lambda}) \subset \mathcal{S}_m(\boldsymbol{\nu}) \subset \mathcal{S}_m(\boldsymbol{\mu})} (-1)^{\#\mathcal{S}_m(\boldsymbol{\nu})+1} = \begin{cases} (-1)^{l(\boldsymbol{\mu})} & \text{if } \boldsymbol{\lambda} = \boldsymbol{\mu} \\ 0 & \text{if } \boldsymbol{\lambda} < \boldsymbol{\mu}. \end{cases}$$

Therefore we obtain $d\sigma d(\mu) = \mu$.

(iii) The proof of (iii) is similar to the proof of (ii).

(iv) We have

$$d * \sigma d(\mu) = \sum_{\nu \leq \mu} (-1)^{l(\nu)} \sum_{\lambda \leq \nu^*} \lambda = \sum_{\lambda \in I_m} \left\{ \sum_{\substack{\nu \leq \mu \\ \nu \leq \lambda^*}} (-1)^{l(\nu)} \right\} \lambda.$$

By Lemma 2.2, we have

$$\sum_{\substack{\nu \leq \mu \\ \nu \leq \lambda^*}} (-1)^{l(\nu)} = \sum_{\mathcal{S}_m(\nu) \subset \mathcal{S}_m(\mu) \cap \mathcal{S}_m(\lambda)^c} (-1)^{\#\mathcal{S}_m(\nu)+1} = \begin{cases} -1 & \text{if } \mathcal{S}_m(\mu) \subset \mathcal{S}_m(\lambda) \\ 0 & \text{if } \mathcal{S}_m(\mu) \not\subset \mathcal{S}_m(\lambda). \end{cases}$$

Hence we get

$$d * \sigma d(\mu) = - \sum_{\lambda \geq \mu} \lambda = -u(\mu).$$

This implies the assertion.

(v) The proof of (v) is similar to the proof of (iv). $\square$

We define the multiplication $*$ on V known as the harmonic product. We shall give a combinatorial description of the harmonic product. For $\mu, \nu \in I$, we consider $2 \times l$ ($l \geq 1$) matrices $M = (m_{ij})$ with the following properties:

- Each entry of M is a non-negative integer.
- If we exclude zeros from the first row of M , we get μ .
- If we exclude zeros from the second row of M , we get ν .
- For $1 \leq j \leq l$, we have $m_{1j} + m_{2j} > 0$.

(Then it must hold that $\max\{l(\mu), l(\nu)\} \leq l \leq l(\mu) + l(\nu)$.) We denote the set of all such matrices by $\mathcal{H}_{\mu, \nu}$. For example, for $\mu = (1)$ and $\nu = (2, 3)$, we have

$$\mathcal{H}_{\mu, \nu} = \left\{ \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 3 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 2 & 0 & 3 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 2 & 3 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 2 & 3 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 2 & 3 \end{pmatrix} \right\}. \quad (5)$$

Definition 2.4. For $\mu, \nu \in I$, we define

$$\mu * \nu = \sum_{M=(m_{ij}) \in \mathcal{H}_{\mu, \nu}} (m_{11} + m_{21}, \dots, m_{1l} + m_{2l}) \in V$$

and

$$\mu \bar{*} \nu = \sum_{M=(m_{ij}) \in \mathcal{H}_{\mu, \nu}} (-1)^{l(\mu)+l(\nu)-l} (m_{11} + m_{21}, \dots, m_{1l} + m_{2l}) \in V,$$

where l is the number of columns of M . We extend $*$, $\bar{*}$: $I \times I \rightarrow V$ bilinearly on $V \times V$.

For example, we have

$$(1) * (2, 3) = (1, 2, 3) + (2, 1, 3) + (2, 3, 1) + (3, 3) + (2, 4)$$

and

$$(1) \overline{*} (2, 3) = (1, 2, 3) + (2, 1, 3) + (2, 3, 1) - (3, 3) - (2, 4)$$

from (5).

Proposition 2.5. *For any $v, w \in V$, we have $d(v \overline{*} w) = d(v) * d(w)$.*

Proof. For a proof of this proposition, see [14, Proof of Proposition 2.4]. $\square$

For $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$, $\boldsymbol{\nu} = (\nu_1, \dots, \nu_q) \in I$, we define

$$\begin{aligned}\boldsymbol{\mu} \# \boldsymbol{\nu} &= (\mu_1, \dots, \mu_p, \nu_1, \dots, \nu_q), \\ \boldsymbol{\mu} \dot{\#} \boldsymbol{\nu} &= (\mu_1, \dots, \mu_{p-1}, \mu_p + \nu_1, \nu_2, \dots, \nu_q)\end{aligned}$$

and

$$\boldsymbol{\mu}^+ = (\mu_1, \dots, \mu_{p-1}, \mu_p + 1).$$

In addition, if $|\boldsymbol{\mu}| > 1$, we define

$$\boldsymbol{\mu}^- = \begin{cases} (\mu_1, \dots, \mu_{p-1}, \mu_p - 1) & \text{if } \mu_p > 1 \\ (\mu_1, \dots, \mu_{p-1}) & \text{if } \mu_p = 1. \end{cases}$$

We extend $\#, \dot{\#}$: $I \times I \rightarrow V$ bilinearly on $V \times V$ and $\boldsymbol{\mu} \mapsto \boldsymbol{\mu}^+$ linearly on V . The following Propositions 2.6 and 2.7 are easily proved.

Proposition 2.6. *For any $\boldsymbol{\mu}, \boldsymbol{\nu} \in I$, we have $(\boldsymbol{\mu} \# \boldsymbol{\nu})^* = \boldsymbol{\mu}^* \dot{\#} \boldsymbol{\nu}^*$.*

Proposition 2.7. *For any $\boldsymbol{\mu}, \boldsymbol{\nu} \in I$, we have*

- (i) $u(\boldsymbol{\mu} \# \boldsymbol{\nu}) = u(\boldsymbol{\mu}) \# u(\boldsymbol{\nu})$,
- (ii) $u(\boldsymbol{\mu} \dot{\#} \boldsymbol{\nu}) = u(\boldsymbol{\mu}) \# u(\boldsymbol{\nu}) + u(\boldsymbol{\mu}) \dot{\#} u(\boldsymbol{\nu})$,
- (iii) $d(\boldsymbol{\mu} \dot{\#} \boldsymbol{\nu}) = d(\boldsymbol{\mu}) \dot{\#} d(\boldsymbol{\nu})$,
- (iv) $d(\boldsymbol{\mu} \# \boldsymbol{\nu}) = d(\boldsymbol{\mu}) \# d(\boldsymbol{\nu}) + d(\boldsymbol{\mu}) \dot{\#} d(\boldsymbol{\nu})$.

Here, we introduce the empty multi-index ϕ . We denote $\{\phi\} \cup I$ by $\tilde{I}$ and $\mathbb{Q}\phi \oplus V$ by $\tilde{V}$. It is natural to define

$$\phi^+ = (1) \quad \text{and} \quad (1)^- = \phi.$$

We define $*(\phi)$, $\tau(\phi)$, $\sigma(\phi)$, $u(\phi)$ and $d(\phi)$ to be ϕ . Moreover, we define for any $\boldsymbol{\mu} \in \tilde{I}$

$$\phi * \boldsymbol{\mu} = \boldsymbol{\mu} * \phi = \boldsymbol{\mu}, \quad \phi \overline{*} \boldsymbol{\mu} = \boldsymbol{\mu} \overline{*} \phi = \boldsymbol{\mu}$$

and

$$\phi \# \boldsymbol{\mu} = \boldsymbol{\mu} \# \phi = \boldsymbol{\mu}, \quad \phi \dot{\#} \boldsymbol{\mu} = \boldsymbol{\mu} \dot{\#} \phi = \boldsymbol{\mu}.$$

Proposition 2.8 follows from the definition of the harmonic product $*$.

Proposition 2.8. For any $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p), \boldsymbol{\nu} = (\nu_1, \dots, \nu_q) \in I$, we have

$$\boldsymbol{\mu} * \boldsymbol{\nu} = (\boldsymbol{\mu}^- * \boldsymbol{\nu}) \# (\mu_p) + (\boldsymbol{\mu} * \boldsymbol{\nu}^-) \# (\nu_q) + (\boldsymbol{\mu}^- * \boldsymbol{\nu}^-) \# (\mu_p + \nu_q).$$

In the last of this section, we define bilinear mappings $\otimes, \overline{\otimes}: V \times V \rightarrow V$, which are used in the following sections, and prove a proposition (Proposition 2.11). Lemma 2.9 is used in the proof of Proposition 2.11. We note that

$$d(\boldsymbol{\mu}) = \sum_{i=1}^p d(\mu_1, \dots, \mu_{i-1}) \# (\mu_i + \dots + \mu_p) \quad (6)$$

for any $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$, which is immediate from the definition of d .

Lemma 2.9. For any $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p), \boldsymbol{\nu} = (\nu_1, \dots, \nu_q) \in I$, we have

$$\begin{aligned} d(\boldsymbol{\mu}) * d(\boldsymbol{\nu}) &= \sum_{i=1}^p \left(d(\mu_1, \dots, \mu_{i-1}) * d(\boldsymbol{\nu}) \right) \# (\mu_i + \dots + \mu_p) \\ &+ \sum_{j=1}^q \left(d(\boldsymbol{\mu}) * d(\nu_1, \dots, \nu_{j-1}) \right) \# (\nu_j + \dots + \nu_q) \\ &+ \sum_{i=1}^p \sum_{j=1}^q \left(d(\mu_1, \dots, \mu_{i-1}) * d(\nu_1, \dots, \nu_{j-1}) \right) \# (\mu_i + \dots + \mu_p + \nu_j + \dots + \nu_q). \end{aligned}$$

Proof. It follows from (6) and Proposition 2.8. $\square$

Definition 2.10. For $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p), \boldsymbol{\nu} = (\nu_1, \dots, \nu_q) \in I$, we define

$$\boldsymbol{\mu} \otimes \boldsymbol{\nu} = (\boldsymbol{\mu}^- * \boldsymbol{\nu}^-) \# (\mu_p + \nu_q)$$

and

$$\boldsymbol{\mu} \overline{\otimes} \boldsymbol{\nu} = (\boldsymbol{\mu}^- \overline{*} \boldsymbol{\nu}^-) \# (\mu_p + \nu_q).$$

We extend $\otimes, \overline{\otimes}: I \times I \rightarrow V$ bilinearly on $V \times V$.

Proposition 2.11. For any $v, w \in V$, we have $d(v \overline{\otimes} w) = d(v) \otimes d(w)$.

Proof. It suffices to show that the equality $d(\boldsymbol{\mu} \overline{\otimes} \boldsymbol{\nu}) = d(\boldsymbol{\mu}) \otimes d(\boldsymbol{\nu})$ holds for any $\boldsymbol{\mu}, \boldsymbol{\nu} \in I$. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p), \boldsymbol{\nu} = (\nu_1, \dots, \nu_q) \in I$. Then we have

$$d(\boldsymbol{\mu} \overline{\otimes} \boldsymbol{\nu}) = \left(d(\boldsymbol{\mu}^-) * d(\boldsymbol{\nu}^-) \right) \# (\mu_p + \nu_q) + \left(d(\boldsymbol{\mu}^-) * d(\boldsymbol{\nu}^-) \right) \# (\mu_p + \nu_q) \quad (7)$$

by the definition of $\overline{\otimes}$, Propositions 2.7 and 2.5. By Lemma 2.9, we have

$$\begin{aligned}
& \left(d(\boldsymbol{\mu}^-) * d(\boldsymbol{\nu}^-) \right) \# (\mu_p + \nu_q) \\
&= \sum_{i=1}^{p-1} \left(d(\mu_1, \dots, \mu_{i-1}) * d(\boldsymbol{\nu}^-) \right) \# (\mu_i + \dots + \mu_p + \nu_q) \\
&+ \sum_{j=1}^{q-1} \left(d(\boldsymbol{\mu}^-) * d(\nu_1, \dots, \nu_{j-1}) \right) \# (\mu_p + \nu_j + \dots + \nu_q) \\
&+ \sum_{i=1}^{p-1} \sum_{j=1}^{q-1} \left(d(\mu_1, \dots, \mu_{i-1}) * d(\nu_1, \dots, \nu_{j-1}) \right) \# (\mu_i + \dots + \mu_p + \nu_j + \dots + \nu_q).
\end{aligned}$$

Hence the right-hand side of (7) equals

$$\sum_{i=1}^p \sum_{j=1}^q \left(d(\mu_1, \dots, \mu_{i-1}) * d(\nu_1, \dots, \nu_{j-1}) \right) \# (\mu_i + \dots + \mu_p + \nu_j + \dots + \nu_q),$$

which is equal to $d(\boldsymbol{\mu}) \otimes d(\boldsymbol{\nu})$ by (6). $\square$

3 A formula for finite multiple harmonic sums

In this section, we prove a formula for finite multiple harmonic sums (Theorem 3.8). We note that $\mathbb{N}$ denotes the set of non-negative integers.

Definition 3.1. (i) We define the difference $\Delta: \mathbb{C}^{\mathbb{N}} \rightarrow \mathbb{C}^{\mathbb{N}}$ by putting

$$(\Delta a)(n) = a(n) - a(n+1)$$

for $a \in \mathbb{C}^{\mathbb{N}}$ and $n \in \mathbb{N}$. We denote the composition of Δ with itself n times by Δ^n .

(ii) We define the inversion $\nabla: \mathbb{C}^{\mathbb{N}} \rightarrow \mathbb{C}^{\mathbb{N}}$ by putting

$$(\nabla a)(n) = (\Delta^n a)(0)$$

for $a \in \mathbb{C}^{\mathbb{N}}$ and $n \in \mathbb{N}$.

Let $\mathbb{C}[[x, y]]$ be the ring of formal power series over the complex numbers in two indeterminates x and y . We put $\partial_x = \partial/\partial x$ and $\partial_y = \partial/\partial y$. For

$$f(x, y) = \sum_{n, k=0}^{\infty} a(n, k) \frac{x^n y^k}{n! k!} \in \mathbb{C}[[x, y]],$$

we have

$$(\partial_x + \partial_y - 1)f(x, y) = \sum_{n, k=0}^{\infty} \{a(n+1, k) + a(n, k+1) - a(n, k)\} \frac{x^n y^k}{n! k!}. \quad (8)$$

Therefore by the definition of the difference we have

$$(\partial_x + \partial_y - 1) \sum_{n,k=0}^{\infty} (\Delta^k a)(n) \frac{x^n y^k}{n! k!} = 0. \quad (9)$$

Lemma 3.2. *Let $f(x, y) \in \mathbb{C}[[x, y]]$. If $f(x, y)$ satisfies two conditions*

$$(\partial_x + \partial_y - 1)f(x, y) = 0 \quad \text{and} \quad f(x, 0) = 0,$$

we have $f(x, y) = 0$.

Proof. Let

$$f(x, y) = \sum_{n,k=0}^{\infty} a(n, k) \frac{x^n y^k}{n! k!}.$$

By (8), the conditions of the lemma are equivalent to

$$\begin{cases} a(n+1, k) + a(n, k+1) = a(n, k) & (n, k \in \mathbb{N}), \\ a(n, 0) = 0 & (n \in \mathbb{N}). \end{cases}$$

By induction on k , we obtain $a(n, k) = 0$ for any $n, k \in \mathbb{N}$. □

Proposition 3.3. *Let $a \in \mathbb{C}^{\mathbb{N}}$. For any $n, k \in \mathbb{N}$, we have*

$$(\Delta^k(\nabla a))(n) = (\Delta^n a)(k).$$

In particular, we obtain $\nabla(\nabla a) = a$ by setting $n = 0$.

Proof. For a sequence $a \in \mathbb{C}^{\mathbb{N}}$, we put

$$f_a(x, y) = \sum_{n,k=0}^{\infty} (\Delta^k a)(n) \frac{x^n y^k}{n! k!}.$$

By (9) we have $(\partial_x + \partial_y - 1)f_{\nabla a}(x, y) = 0$ and $(\partial_x + \partial_y - 1)f_a(y, x) = 0$. We also have

$$f_{\nabla a}(x, 0) = \sum_{n=0}^{\infty} (\nabla a)(n) \frac{x^n}{n!} = f_a(0, x)$$

by the definition of the inversion ∇ . Therefore by Lemma 3.2, we obtain $f_{\nabla a}(x, y) = f_a(y, x)$. This completes the proof of Proposition 3.3. □

Let

$$F(x) = \sum_{n=0}^{\infty} a(n) \frac{x^n}{n!}.$$

By using Lemma 3.2, we can easily prove that

$$F(x - y)e^y = \sum_{n,k=0}^{\infty} (\Delta^k a)(n) \frac{x^n y^k}{n! k!}.$$

By comparing the coefficients of $x^n y^k$ on both sides, we obtain the following proposition.

Proposition 3.4. For any $n, k \in \mathbb{N}$, we have

$$(\Delta^k a)(n) = \sum_{i=0}^k (-1)^i \binom{k}{i} a(n+i).$$

In particular, we obtain

$$(\nabla a)(k) = \sum_{i=0}^k (-1)^i \binom{k}{i} a(i)$$

for any $k \in \mathbb{N}$ by setting $n = 0$.

Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$ and $n \in \mathbb{N}$. In this paper, we consider four kinds of finite multiple harmonic sums

$$\begin{aligned} s_{\boldsymbol{\mu}}(n) &= \sum_{0 \leq n_1 \leq \dots \leq n_p = n} \frac{1}{(n_1 + 1)^{\mu_1} \dots (n_p + 1)^{\mu_p}}, \\ a_{\boldsymbol{\mu}}(n) &= \sum_{0 \leq n_1 < \dots < n_p = n} \frac{1}{(n_1 + 1)^{\mu_1} \dots (n_p + 1)^{\mu_p}}, \\ S_{\boldsymbol{\mu}}(n) &= \sum_{0 \leq n_1 \leq \dots \leq n_p < n} \frac{1}{(n_1 + 1)^{\mu_1} \dots (n_p + 1)^{\mu_p}} \end{aligned}$$

and

$$A_{\boldsymbol{\mu}}(n) = \sum_{0 \leq n_1 < \dots < n_p < n} \frac{1}{(n_1 + 1)^{\mu_1} \dots (n_p + 1)^{\mu_p}}.$$

We extend the mappings $I \ni \boldsymbol{\mu} \mapsto s_{\boldsymbol{\mu}}, a_{\boldsymbol{\mu}}, S_{\boldsymbol{\mu}}, A_{\boldsymbol{\mu}} \in \mathbb{C}^{\mathbb{N}}$ linearly on V . We introduced the empty multi-index ϕ in Section 2. We define

$$S_{\phi}(n) = A_{\phi}(n) = 1$$

for any $n \in \mathbb{N}$.

Proposition 3.5. (i) For any $\boldsymbol{\mu} \in I$, we have $-(\Delta S_{\boldsymbol{\mu}}) = s_{\boldsymbol{\mu}}$ and $-(\Delta A_{\boldsymbol{\mu}}) = a_{\boldsymbol{\mu}}$.
(ii) For any $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$ and any $n \in \mathbb{N}$, we have

$$s_{\boldsymbol{\mu}}(n) = S_{\mu_1, \dots, \mu_{p-1}}(n+1) \frac{1}{(n+1)^{\mu_p}} \quad \text{and} \quad a_{\boldsymbol{\mu}}(n) = A_{\mu_1, \dots, \mu_{p-1}}(n) \frac{1}{(n+1)^{\mu_p}},$$

where we put $(\mu_1, \dots, \mu_{p-1}) = \phi$ if $p = 1$.

Proof. These are easily seen. $\square$

Proposition 3.6. (i) For any $v \in V$, we have $s_v = a_{d(v)}$, $S_v = A_{d(v)}$, $a_v = s_{d^{-1}(v)}$ and $A_v = S_{d^{-1}(v)}$.
(ii) For any $v, w \in V$, we have $A_v A_w = A_{v * w}$, $S_v S_w = S_{v \overline{*} w}$, $a_v a_w = a_{v \otimes w}$ and $s_v s_w = s_{v \overline{\otimes} w}$.

Proof. (i) It is easily seen that $s_{\mu} = a_{d(\mu)}$ and $S_{\mu} = A_{d(\mu)}$ for $\mu \in I$. The assertions of (i) follow immediately from these.

(ii) It is easily seen that $A_{\mu}A_{\nu} = A_{\mu*\nu}$ for any $\mu, \nu \in I$. Therefore the first assertion follows. In addition, for any $v, w \in V$ we have

$$S_v S_w = A_{d(v)} A_{d(w)} = A_{d(v)*d(w)} = S_{d^{-1}(d(v)*d(w))} = S_{v\bar{*}w}$$

by Proposition 2.5. The other assertions follow from Proposition 3.5 (ii) and the definitions of $\otimes$ and $\bar{\otimes}$. $\square$

Let m be a positive integer. For $\mu = (\mu_1, \dots, \mu_p), \nu = (\nu_1, \dots, \nu_q) \in I_m$ and $n, k \in \mathbb{N}$, we also consider a finite sum

$$s_{\mu, \nu}(n, k) = \binom{n+k}{n}^{-1} \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_q = k}} \frac{1}{(n_{i_1} + k_{j_1} + 1) \cdots (n_{i_m} + k_{j_m} + 1)},$$

where

$$(i_1, \dots, i_m) = (\underbrace{1, \dots, 1}_{\mu_1}, \dots, \underbrace{p, \dots, p}_{\mu_p})$$

and

$$(j_1, \dots, j_m) = (\underbrace{1, \dots, 1}_{\nu_1}, \dots, \underbrace{q, \dots, q}_{\nu_q}).$$

We note that

$$s_{\mu, \nu}(n, 0) = s_{\mu}(n) \quad \text{and} \quad s_{\mu, \nu}(0, k) = s_{\nu}(k). \quad (10)$$

Lemma 3.7. *Let $\mu = (\mu_1, \dots, \mu_p), \nu = (\nu_1, \dots, \nu_q) \in I$ and let $|\mu| = |\nu| = m$. Then for any $n, k \in \mathbb{N}$, we have*

$$s_{\mu^-, \nu^-}(n, k) = \begin{cases} (n+k+1)s_{\mu, \nu}(n, k) - k s_{\mu, \nu}(n, k-1) & \text{if } \mu_p > 1 \text{ and } \nu_q = 1 \\ (n+k+1)s_{\mu, \nu}(n, k) - n s_{\mu, \nu}(n-1, k) & \text{if } \mu_p = 1 \text{ and } \nu_q > 1, \end{cases}$$

where we put $k s_{\mu, \nu}(n, k-1) = 0$ if $k = 0$ and $n s_{\mu, \nu}(n-1, k) = 0$ if $n = 0$.

Proof. We shall prove the lemma only for $\mu_p > 1$ and $\nu_q = 1$. We can prove the lemma for $\mu_p = 1$ and $\nu_q > 1$ similarly. We have

$$\begin{aligned} & (n+k+1)s_{\mu, \nu}(n, k) - k s_{\mu, \nu}(n, k-1) \\ &= \binom{n+k}{n}^{-1} (n+k+1) \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_q = k}} \frac{1}{(n_{i_1} + k_{j_1} + 1) \cdots (n_{i_m} + k_{j_m} + 1)} \\ & \quad - \binom{n+k}{n}^{-1} (n+k) \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_q = k-1}} \frac{1}{(n_{i_1} + k_{j_1} + 1) \cdots (n_{i_m} + k_{j_m} + 1)}. \end{aligned}$$

Since $\mu_p > 1$ and $\nu_q = 1$, this equals

$$\begin{aligned} & \binom{n+k}{n}^{-1} \left\{ \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_{q-1} \leq k}} \frac{1}{(n_{i_1} + k_{j_1} + 1) \cdots (n_{i_{m-1}} + k_{j_{m-1}} + 1)} \right. \\ & \quad \left. - \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_{q-1} \leq k-1}} \frac{1}{(n_{i_1} + k_{j_1} + 1) \cdots (n_{i_{m-1}} + k_{j_{m-1}} + 1)} \right\} \\ & = \binom{n+k}{n}^{-1} \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_{q-1} = k}} \frac{1}{(n_{i_1} + k_{j_1} + 1) \cdots (n_{i_{m-1}} + k_{j_{m-1}} + 1)}. \end{aligned}$$

We have thus proved the lemma. $\square$

Theorem 3.8. For any $\mu \in I$ and $n, k \in \mathbb{N}$ we have $(\Delta^k s_\mu)(n) = s_{\mu, \mu^*}(n, k)$.

Proof. For $\mu \in I$, we put

$$f_\mu(x, y) = \sum_{n, k=0}^{\infty} (\Delta^k s_\mu)(n) \frac{x^n y^k}{n! k!}$$

and for $\mu, \nu \in I$ with $|\mu| = |\nu|$, we put

$$g_{\mu, \nu}(x, y) = \sum_{n, k=0}^{\infty} s_{\mu, \nu}(n, k) \frac{x^n y^k}{n! k!}.$$

If we have

$$(\partial_x + \partial_y - 1)g_{\mu, \mu^*}(x, y) = 0, \quad (11)$$

then by Lemma 3.2 and (10) we have $f_\mu(x, y) = g_{\mu, \mu^*}(x, y)$. Therefore we have only to prove (11). If we put

$$\xi = x\partial_x + y\partial_y + 1 - y \quad \text{and} \quad \eta = x\partial_x + y\partial_y + 1 - x,$$

by Lemma 3.7 we have

$$g_{\mu^-, \nu^-} = \begin{cases} \xi g_{\mu, \nu} & \text{if } \mu_p > 1 \text{ and } \nu_q = 1 \\ \eta g_{\mu, \nu} & \text{if } \mu_p = 1 \text{ and } \nu_q > 1. \end{cases}$$

We note that

$$(\partial_x + \partial_y - 1)\xi - \xi(\partial_x + \partial_y + 1) = \partial_x + \partial_y - 1$$

and

$$(\partial_x + \partial_y - 1)\eta - \eta(\partial_x + \partial_y + 1) = \partial_x + \partial_y - 1.$$

We prove (11) by induction on $|\boldsymbol{\mu}|$. For $|\boldsymbol{\mu}| = 1$ (i.e. $\boldsymbol{\mu} = (1)$), we have

$$g_{\boldsymbol{\mu}, \boldsymbol{\mu}^*}(x, y) = \sum_{n, k=0}^{\infty} \frac{x^n y^k}{(n+k+1)!}.$$

Therefore we can easily check that (11) holds in this case. We assume that the claim (11) is true for $|\boldsymbol{\mu}| = m$. Let $|\boldsymbol{\mu}| = m+1$, $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$ and $\boldsymbol{\mu}^* = (\mu_1^*, \dots, \mu_q^*)$. If $\mu_p > 1$, then we have $\mu_q^* = 1$ by (4). Therefore we obtain

$$\begin{aligned} (\xi + 1)(\partial_x + \partial_y - 1)g_{\boldsymbol{\mu}, \boldsymbol{\mu}^*} &= (\partial_x + \partial_y - 1)\xi g_{\boldsymbol{\mu}, \boldsymbol{\mu}^*} = (\partial_x + \partial_y - 1)g_{\boldsymbol{\mu}^-, (\boldsymbol{\mu}^*)^-} \\ &= (\partial_x + \partial_y - 1)g_{\boldsymbol{\mu}^-, (\boldsymbol{\mu}^-)^*} = 0 \end{aligned}$$

by the hypothesis of the induction. If $\mu_p = 1$, then we obtain $(\eta + 1)(\partial_x + \partial_y - 1)g_{\boldsymbol{\mu}, \boldsymbol{\mu}^*} = 0$ by a similar argument. We can easily check that mappings $\xi + 1$, $\eta + 1: \mathbb{C}[[x, y]] \rightarrow \mathbb{C}[[x, y]]$ are injections. Thus we have proved (11). $\square$

Corollary 3.9. *For any $\boldsymbol{\mu} \in I$, we have $\nabla s_{\boldsymbol{\mu}} = s_{\boldsymbol{\mu}^*}$.*

Proof. If we set $n = 0$ in Theorem 3.8, we obtain the result by (10). $\square$

We define $T: \mathbb{C}^{\mathbb{N}} \rightarrow \mathbb{C}^{\mathbb{N}}$ by

$$(Ta)(n) = a(n+1) \quad (a \in \mathbb{C}^{\mathbb{N}}, n \in \mathbb{N}),$$

and then we have $\Delta = \nabla T \nabla$.

Corollary 3.10. *For any $\boldsymbol{\mu} \in I$ and $n \in \mathbb{N}$, we have*

$$(\nabla S_{\boldsymbol{\mu}})(n) = \begin{cases} 0 & \text{if } n = 0 \\ -s_{\boldsymbol{\mu}^*}(n-1) & \text{if } n \geq 1. \end{cases}$$

Proof. By Proposition 3.5 (i) and Corollary 3.9 we have $T \nabla S_{\boldsymbol{\mu}} = -s_{\boldsymbol{\mu}^*}$, which completes the proof. $\square$

The following estimation is used in Section 5.

Proposition 3.11. *Let $k \in \mathbb{N}$ and let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$. For any $\varepsilon > 0$, we have*

$$(\Delta^k s_{\boldsymbol{\mu}})(n) = O\left(\frac{1}{n^{k+\mu_p-\varepsilon}}\right) \quad (n \rightarrow \infty).$$

Proof. Let $|\boldsymbol{\mu}| = m$ and let $\boldsymbol{\mu}^* = (\mu_1^*, \dots, \mu_q^*)$. By Theorem 3.8 we have

$$(\Delta^k s_{\boldsymbol{\mu}})(n) = \binom{n+k}{n}^{-1} \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_q = k}} \frac{1}{(n_{i_1} + k_{j_1} + 1) \cdots (n_{i_m} + k_{j_m} + 1)},$$

where

$$(i_1, \dots, i_m) = (\underbrace{1, \dots, 1}_{\mu_1}, \dots, \underbrace{p, \dots, p}_{\mu_p}), \quad (j_1, \dots, j_m) = (\underbrace{1, \dots, 1}_{\mu_1^*}, \dots, \underbrace{q, \dots, q}_{\mu_q^*}).$$

The number of elements of the set

$$\{(k_1, \dots, k_{q-1}) \in \mathbb{N}^{q-1} \mid 0 \leq k_1 \leq \dots \leq k_{q-1} \leq k\}$$

is bounded by $(k+1)^{q-1}$. Hence we have

$$\begin{aligned} & \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_q = k}} \frac{1}{(n_{i_1} + k_{j_1} + 1) \cdots (n_{i_m} + k_{j_m} + 1)} \\ & \leq \sum_{\substack{0 \leq n_1 \leq \dots \leq n_p = n \\ 0 \leq k_1 \leq \dots \leq k_q = k}} \frac{1}{(n_1 + 1)^{\mu_1} \cdots (n_p + 1)^{\mu_p}} \\ & \leq (k+1)^{q-1} \sum_{0 \leq n_1 \leq \dots \leq n_p = n} \frac{1}{(n_1 + 1)^{\mu_1} \cdots (n_p + 1)^{\mu_p}} \\ & \leq \frac{(k+1)^{q-1}}{(n+1)^{\mu_p}} \left(\sum_{0 \leq n_1 \leq n} \frac{1}{n_1 + 1} \right) \cdots \left(\sum_{0 \leq n_{p-1} \leq n} \frac{1}{n_{p-1} + 1} \right) \\ & = O\left(\frac{\log^{p-1} n}{n^{\mu_p}}\right) \quad (n \rightarrow \infty), \end{aligned}$$

which completes the proof. $\square$

Moreover, we use the inequality

$$s_{\mu_1, \dots, \mu_p}(n) \geq \frac{1}{(n+1)^{\mu_p}} \quad (n \in \mathbb{N}) \quad (12)$$

in Section 5.

4 The Newton series

In this section, we state general properties of the Newton series needed later. We do not prove Propositions 4.1, 4.2 and 4.4. For proofs of these propositions, see [3, 9, 13]. For a sequence $a \in \mathbb{C}^{\mathbb{N}}$, the series with complex variable z

$$f(z) = \sum_{n=0}^{\infty} (-1)^n (\nabla a)(n) \binom{z}{n}, \quad \binom{z}{n} = \frac{z(z-1) \cdots (z-n+1)}{n!} \quad (13)$$

is called the Newton series which interpolates a sequence a . In fact, for $z = n \in \mathbb{N}$, the sum of this series is

$$\sum_{k=0}^n (-1)^k (\nabla a)(k) \binom{n}{k} = (\nabla^2 a)(n) = a(n)$$

by Propositions 3.4 and 3.3. We first state known results for the convergence of the Newton series.

Proposition 4.1. *Let $a \in \mathbb{C}^{\mathbb{N}}$ and $z_0 \in \mathbb{C} \setminus \mathbb{N}$. If the Newton series*

$$\sum_{n=0}^{\infty} (-1)^n a(n) \binom{z}{n}$$

converges at $z = z_0$, then the series converges uniformly in the wider sense in the region $\operatorname{Re} z > \operatorname{Re} z_0$.

By Proposition 4.1, there exists the unique $\rho \in \mathbb{R} \cup \{\pm\infty\}$ such that the series $f(z)$ defined by (13) converges if $\operatorname{Re} z > \rho$ and does not converge if $\operatorname{Re} z < \rho$ and $z \notin \mathbb{N}$. We call ρ the abscissa of convergence of the series $f(z)$. The function $f(z)$ is analytic in the region $\operatorname{Re} z > \rho$.

Proposition 4.2. *Let $a \in \mathbb{C}^{\mathbb{N}}$ and $z \in \mathbb{C} \setminus \mathbb{N}$. Then the Newton series*

$$\sum_{n=0}^{\infty} (-1)^n a(n) \binom{z}{n}$$

converges (resp. converges absolutely) if and only if the Dirichlet series

$$\sum_{n=0}^{\infty} \frac{a(n)}{(n+1)^{z+1}}$$

converges (resp. converges absolutely).

The following corollary is immediate from Proposition 4.2.

Corollary 4.3. *Let $a \in \mathbb{C}^{\mathbb{N}}$ and $\varepsilon \in \mathbb{R}$. If we have $a(n) = O(n^\varepsilon)$ ($n \rightarrow \infty$), then the Newton series*

$$\sum_{n=0}^{\infty} (-1)^n a(n) \binom{z}{n}$$

converges absolutely for any $z \in \mathbb{C}$ with $\operatorname{Re} z > \varepsilon$.

The function $f(z)$ defined by a Newton series is uniquely determined by $f(n)$ for all sufficiently large integers n .

Proposition 4.4. *Let $a \in \mathbb{C}^{\mathbb{N}}$ and let the Newton series*

$$f(z) = \sum_{n=0}^{\infty} (-1)^n a(n) \binom{z}{n}$$

have the abscissa of convergence ρ . If there exists $N \in \mathbb{N}$ such that $f(n) = 0$ for any integers $n \geq N$, then we have $f(z) = 0$ for any $\operatorname{Re} z > \rho$.

The product of two Newton series does not necessarily have a Newton series expansion. We give a sufficient condition for the product of two Newton series to be expanded to a Newton series. We need a lemma.

Lemma 4.5. *Let $a \in \mathbb{C}^{\mathbb{N}}$ and let the Newton series*

$$f(z) = \sum_{n=0}^{\infty} (-1)^n a(n) \binom{z}{n}$$

have the abscissa of convergence ρ . Let $k \in \mathbb{N}$. Then for any $\operatorname{Re} z > \rho + k$, we have

$$(-1)^k \binom{z}{k} f(z) = \sum_{n=k}^{\infty} (-1)^n \binom{n}{k} (\Delta^k a)(n-k) \binom{z}{n}.$$

Proof. For any $z \in \mathbb{C}$ and $n, k \in \mathbb{N}$, we have

$$\binom{z}{n} = \sum_{i=0}^k \binom{k}{i} \binom{z-k}{n-i}.$$

Hence we have for any $\operatorname{Re} z > \rho$

$$\begin{aligned} & (-1)^k \binom{z}{k} f(z) \\ &= (-1)^k \sum_{i=0}^k \binom{k}{i} \sum_{n=i}^{\infty} (-1)^n a(n) \binom{z}{k} \binom{z-k}{n-i} \\ &= (-1)^k \sum_{i=0}^k \binom{k}{i} \sum_{n=i}^{\infty} (-1)^n a(n) \binom{k+n-i}{k} \binom{z}{k+n-i} \\ &= (-1)^k \sum_{i=0}^k \binom{k}{i} \sum_{n=0}^{\infty} (-1)^{n+i} a(n+i) \binom{k+n}{k} \binom{z}{k+n}. \end{aligned}$$

By Proposition 3.4, this is equal to

$$(-1)^k \sum_{n=0}^{\infty} (-1)^n (\Delta^k a)(n) \binom{k+n}{k} \binom{z}{k+n},$$

which completes the proof. $\square$

Proposition 4.6. *Let $a, b \in \mathbb{C}^{\mathbb{N}}$. Let the Newton series*

$$f(z) = \sum_{n=0}^{\infty} (-1)^n a(n) \binom{z}{n} \quad \text{and} \quad g(z) = \sum_{n=0}^{\infty} (-1)^n b(n) \binom{z}{n}$$

have the abscissas of convergence ρ_a and ρ_b , respectively. Let $\varepsilon > 0$. We assume that sequences a and b satisfy the following conditions:

- (i) *The values $a(n)$ and $(\Delta^k b)(n)$ are non-negative for any $n, k \in \mathbb{N}$.*
- (ii) *$\rho_a < 0$.*
- (iii) *For any $k \in \mathbb{N}$ we have $(\Delta^k b)(n) = O(n^{-k-\varepsilon})$ ($n \rightarrow \infty$).*

Then the product $f(z)g(z)$ is expressed by a Newton series in the half-plane $\operatorname{Re} z > \max\{\rho_a, -\varepsilon\}$.

Proof. Let $k \in \mathbb{N}$. By Lemma 4.5, it holds that

$$(-1)^k \binom{z}{k} g(z) = \sum_{n=k}^{\infty} (-1)^n \binom{n}{k} (\Delta^k b)(n-k) \binom{z}{n} \quad (14)$$

for any $\operatorname{Re} z > \rho_b + k$. By setting $k = 0$ in the condition (iii), we have $b(n) = O(n^{-\varepsilon})$ ($n \rightarrow \infty$). According to Corollary 4.3, the Newton series $g(z)$ converges in $\operatorname{Re} z > -\varepsilon$. Again by the condition (iii), we have

$$\binom{n}{k} (\Delta^k b)(n-k) = O(n^{-\varepsilon}) \quad (n \rightarrow \infty).$$

So the right-hand side of (14) converges in $\operatorname{Re} z > -\varepsilon$. Thus we see that the equality (14) is valid for any $\operatorname{Re} z > -\varepsilon$. Let $\operatorname{Re} z > \max\{\rho_a, -\varepsilon\}$. We multiply the both sides of (14) by $a(k)$ and take the sum over all values of k from 0 to ∞ . Consequently, we obtain

$$f(z)g(z) = \sum_{k=0}^{\infty} \sum_{n=k}^{\infty} a(k) (-1)^n \binom{n}{k} (\Delta^k b)(n-k) \binom{z}{n}. \quad (15)$$

If we take z as a negative real number with $z > \max\{\rho_a, -\varepsilon\}$, then each term of the right-hand side of (15) is non-negative from the condition (i). Hence the right-hand side of (15) is equal to

$$\sum_{n=0}^{\infty} (-1)^n \left\{ \sum_{k=0}^n a(k) \binom{n}{k} (\Delta^k b)(n-k) \right\} \binom{z}{n}.$$

Thus we have completed the proof. $\square$

The Taylor expansion of a Newton series is given by the following proposition. We note that the definition of the sequence $a_{\boldsymbol{\mu}}$ ($\boldsymbol{\mu} \in I$) is given in Section 3.

Proposition 4.7. *Let $a \in \mathbb{C}^{\mathbb{N}}$ be any sequence. We suppose that the abscissa of convergence of the Newton series*

$$f(z) = \sum_{n=0}^{\infty} (-1)^n a(n) \binom{z}{n}$$

is negative. Then we have

$$f(z) = a(0) + \sum_{m=1}^{\infty} (-1)^m \left\{ \sum_{n=1}^{\infty} a(n) \underbrace{a_1, \dots, 1}_m (n-1) \right\} z^m$$

in some neighborhood of 0.

Proof. Let $m, n \geq 1$. Then we have

$$\begin{aligned} \left. \frac{1}{m!} \frac{d^m}{dz^m} \binom{z}{n} \right|_{z=0} &= \binom{z}{n} \sum_{0 \leq n_1 < \dots < n_m < n} \frac{1}{(z - n_1) \cdots (z - n_m)} \Big|_{z=0} \\ &= (-1)^{n+m} \underbrace{a_1, \dots, 1}_m (n-1), \end{aligned}$$

which completes the proof. $\square$

5 A class of relations among multiple zeta values

In this section, we consider the Newton series which interpolates the sequence S_μ and derive a class of relations among MZV's. For $\mu \in I$, we define

$$f_\mu(z) = \sum_{n=0}^{\infty} (-1)^n (\nabla s_\mu)(n) \binom{z}{n} \quad \text{and} \quad F_\mu(z) = \sum_{n=0}^{\infty} (-1)^n (\nabla S_\mu)(n) \binom{z}{n}.$$

The abscissas of convergence of $f_\mu(z)$ and $F_\mu(z)$ are given by Proposition 5.1.

Proposition 5.1. *Let $\mu \in I$ and let*

$$\mu = (\mu_1, \dots, \mu_r, \underbrace{1, \dots, 1}_l) \quad (\mu_r \geq 2, r \geq 0, l \geq 0).$$

Then the abscissas of convergence of the Newton series $f_\mu(z)$ and $F_\mu(z)$ are both $-l-1$.

Proof. We prove only the case $F_\mu(z)$. The case $f_\mu(z)$ is proved similarly. Let $\mu^* = (\mu_1^*, \dots, \mu_q^*)$. Then we have $\mu_q^* = l+1$ by (4). By Corollary 3.10 and Proposition 3.11, we obtain

$$(\nabla S_\mu)(n) = O\left(\frac{1}{n^{l+1-\varepsilon}}\right) \quad (n \rightarrow \infty)$$

for any $\varepsilon > 0$. Therefore $F_\mu(z)$ converges for any $\operatorname{Re} z > -l-1$ by Corollary 4.3. Since we have

$$s_{\mu^*}(n) \geq \frac{1}{(n+1)^{l+1}} \quad (n \in N)$$

by (12), the Dirichlet series

$$\sum_{n=0}^{\infty} \frac{(\nabla S_\mu)(n)}{(n+1)^{z+1}} = - \sum_{n=1}^{\infty} \frac{s_{\mu^*}(n-1)}{(n+1)^{z+1}}$$

diverges at $z = -l-1$. Hence the assertion follows from Proposition 4.2. $\square$

More generally, we define

$$F_v(z) = \sum_{n=0}^{\infty} (-1)^n (\nabla S_v)(n) \binom{z}{n}$$

for any $v \in V$. By Proposition 5.1, $F_v(z)$ converges at least in the region $\operatorname{Re} z > -1$ and therefore it can be expanded in a Taylor series at $z = 0$. We shall prove that the Taylor coefficients of $F_v(z)$ is expressed by MZV's. We denote the $\mathbb{Q}$ -vector space whose basis is $I^+ := \{\boldsymbol{\mu}^+ \mid \boldsymbol{\mu} \in I\}$ by V^+ . We define $\mathbb{Q}$ -linear mappings ζ and $\bar{\zeta}$ from V^+ to $\mathbb{R}$ by

$$\zeta(v) = \sum_{n=0}^{\infty} a_v(n) \quad (v \in V^+) \quad \text{and} \quad \bar{\zeta}(v) = \sum_{n=0}^{\infty} s_v(n) \quad (v \in V^+),$$

respectively. It is easily seen that $\bar{\zeta}(v) = \zeta(d(v))$ and $\zeta(v) = \bar{\zeta}(d^{-1}(v))$ from Proposition 3.6 (i).

Proposition 5.2. *Let $v \in V$. Then we have*

$$F_v(z) = \sum_{m=1}^{\infty} (-1)^{m-1} \zeta(d^*(v) \circledast \underbrace{(1, \dots, 1)}_m) z^m$$

in some neighborhood of 0.

Proof. By Proposition 4.7 and Corollary 3.10, we have

$$F_v(z) = \sum_{m=1}^{\infty} (-1)^{m-1} \left\{ \sum_{n=1}^{\infty} s_{v^*}(n-1) \underbrace{a_{1, \dots, 1}}_m(n-1) \right\} z^m.$$

Since

$$\sum_{n=0}^{\infty} s_{v^*}(n) \underbrace{a_{1, \dots, 1}}_m(n) = \sum_{n=0}^{\infty} a_{d^*(v) \circledast \underbrace{(1, \dots, 1)}_m}(n) = \zeta(d^*(v) \circledast \underbrace{(1, \dots, 1)}_m)$$

by Proposition 3.6, the proposition is proved. $\square$

In Section 3, we stated the harmonic product relations for the finite multiple harmonic sums $S_v(n)$ (Proposition 3.6 (ii)). The functions $F_v(z)$ also satisfy the harmonic product relations which come from those for $S_v(n)$.

Theorem 5.3. *Let $v, w \in V$. Then we have*

$$F_v(z)F_w(z) = F_{v \bar{*} w}(z) \tag{16}$$

for any $z \in \mathbb{C}$ with $\operatorname{Re} z > -1$.

Proof. Let $\mu, \nu \in I$. By Lemma 4.5, the function $(z+1)f_{\mu \# (1)}(z)$ is expanded to a Newton series. Therefore we have

$$(z+1)f_{\mu \# (1)}(z) = F_{\mu}(z+1) \quad (\operatorname{Re} z > -2)$$

by Propositions 3.5 (ii) and 4.4. Since $f_{\mu \# (1)}(z)f_{\nu \# (1)}(z)$ is expanded to a Newton series by Propositions 3.11 and 4.6, the product $F_{\mu}(z+1)F_{\nu}(z+1)$ is also expanded to a Newton series. Hence we have

$$F_{\mu}(z+1)F_{\nu}(z+1) = F_{\mu * \nu}(z+1) \quad (\operatorname{Re} z > -2),$$

which completes the proof. $\square$

Now we are in a stage to get the relations among MZV's.

Corollary 5.4. *For any $v, w \in V$ and any integer $m \geq 1$, we have*

$$\sum_{\substack{k+l=m \\ k, l \geq 1}} \zeta(u\sigma(v) \otimes \underbrace{(1, \dots, 1)}_k) \zeta(u\sigma(w) \otimes \underbrace{(1, \dots, 1)}_l) = \zeta(u\sigma(v * w) \otimes \underbrace{(1, \dots, 1)}_m).$$

Proof. We compare the Taylor coefficients in (16). By Proposition 5.2, we have

$$\begin{aligned} \sum_{\substack{k+l=m \\ k, l \geq 1}} \zeta(d*(v) \otimes \underbrace{(1, \dots, 1)}_k) \zeta(d*(w) \otimes \underbrace{(1, \dots, 1)}_l) \\ = -\zeta(d*(v \bar{*} w) \otimes \underbrace{(1, \dots, 1)}_m). \end{aligned}$$

If we replace v and w by $d^{-1}(v)$ and $d^{-1}(w)$, respectively, we obtain the result since we have $d^{-1}(v) \bar{*} d^{-1}(w) = d^{-1}(v * w)$ and $d * d^{-1} = -u\sigma$ by Propositions 2.5 and 2.3. $\square$

We define

$$\zeta^+(v) = \zeta(v^+) \quad \text{and} \quad \bar{\zeta}^+(v) = \bar{\zeta}(v^+)$$

for any $v \in V$. Then it is easily seen that

$$\bar{\zeta}^+(v) = \zeta^+(d(v)). \quad (17)$$

We denote the $\mathbb{Q}$ -vector space generated by $v * w$ for all $v, w \in V$ by $V * V$. The following corollary follows from Corollary 5.4 on setting $m = 1$.

Corollary 5.5. *We have $u\sigma(V * V) \subset \ker \zeta^+$.*

Remark 5.6. We can state Corollary 5.5 as a claim for $\bar{\zeta}^+$. By (17) we have $\bar{\zeta}^+(d^{-1}u\sigma(v * w)) = 0$ for any $v, w \in V$. If we replace v by $d(v)$ and w by $d(w)$, then we obtain

$$\bar{\zeta}^+((v \bar{*} w)^*) = 0$$

from Propositions 2.5 and 2.3.

Remark 5.7. This remark was suggested by the referee. By a calculation similar to

$$A_2(n) = \sum_{0 \leq n_1 < n} \int_0^1 \int_0^1 (x_1 x_2)^{n_1} dx_1 dx_2 = \int_0^1 \int_0^1 \frac{1 - (x_1 x_2)^n}{1 - x_1 x_2} dx_1 dx_2,$$

for any $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$ and $n \in \mathbb{N}$ we obtain

$$A_{\boldsymbol{\mu}}(n) = \int_0^1 \cdots \int_0^1 \sum_{\varepsilon_1, \dots, \varepsilon_{p-1}=0,1} (-1)^{\varepsilon_1 + \cdots + \varepsilon_{p-1}} \frac{1 - (y_1^{\varepsilon_1 \cdots \varepsilon_{p-1}} y_2^{\varepsilon_2 \cdots \varepsilon_{p-1}} \cdots y_{p-1}^{\varepsilon_{p-1}} y_p)^n}{(1 - y_1)(1 - y_1^{\varepsilon_1} y_2) \cdots (1 - y_1^{\varepsilon_1 \cdots \varepsilon_{p-1}} y_2^{\varepsilon_2 \cdots \varepsilon_{p-1}} \cdots y_{p-1}^{\varepsilon_{p-1}} y_p)} dx_1 \cdots dx_{|\boldsymbol{\mu}|},$$

where $y_i = x_{\mu_1 + \cdots + \mu_{i-1} + 1} \cdots x_{\mu_1 + \cdots + \mu_{i-1} + \mu_i}$ ($1 \leq i \leq p$). Therefore we conjecture that

$$G_{\boldsymbol{\mu}}(z) := \sum_{n=0}^{\infty} (-1)^n (\nabla A_{\boldsymbol{\mu}})(n) \binom{z}{n} = \int_0^1 \cdots \int_0^1 \sum_{\varepsilon_1, \dots, \varepsilon_{p-1}=0,1} (-1)^{\varepsilon_1 + \cdots + \varepsilon_{p-1}} \frac{1 - (y_1^{\varepsilon_1 \cdots \varepsilon_{p-1}} y_2^{\varepsilon_2 \cdots \varepsilon_{p-1}} \cdots y_{p-1}^{\varepsilon_{p-1}} y_p)^z}{(1 - y_1)(1 - y_1^{\varepsilon_1} y_2) \cdots (1 - y_1^{\varepsilon_1 \cdots \varepsilon_{p-1}} y_2^{\varepsilon_2 \cdots \varepsilon_{p-1}} \cdots y_{p-1}^{\varepsilon_{p-1}} y_p)} dx_1 \cdots dx_{|\boldsymbol{\mu}|}$$

for $\operatorname{Re} z > 0$. Since

$$G_{\boldsymbol{\mu}}(z) = \sum_{m=1}^{\infty} (-1)^m \zeta(u\sigma(\boldsymbol{\mu}) \otimes \underbrace{(1, \dots, 1)}_m) z^m,$$

we obtain

$$\zeta^+(u\sigma(\boldsymbol{\mu})) = \int_0^1 \cdots \int_0^1 \sum_{\varepsilon_1, \dots, \varepsilon_{p-1}=0,1} (-1)^{\varepsilon_1 + \cdots + \varepsilon_{p-1}} \frac{\log(y_1^{\varepsilon_1 \cdots \varepsilon_{p-1}} y_2^{\varepsilon_2 \cdots \varepsilon_{p-1}} \cdots y_{p-1}^{\varepsilon_{p-1}} y_p)}{(1 - y_1)(1 - y_1^{\varepsilon_1} y_2) \cdots (1 - y_1^{\varepsilon_1 \cdots \varepsilon_{p-1}} y_2^{\varepsilon_2 \cdots \varepsilon_{p-1}} \cdots y_{p-1}^{\varepsilon_{p-1}} y_p)} dx_1 \cdots dx_{|\boldsymbol{\mu}|}.$$

It seems to be possible to prove Corollary 5.5 by standard manipulation on integrals. For example, if $\boldsymbol{\mu} = \boldsymbol{\nu} = (1)$, we have

$$\begin{aligned} \zeta^+(u\sigma(\boldsymbol{\mu} * \boldsymbol{\nu})) &= \zeta^+(u\sigma(2)) + 2\zeta^+(u\sigma(1, 1)) \\ &= \int_0^1 \int_0^1 \left\{ \frac{\log(x_1 x_2)}{1 - x_1 x_2} + \frac{2 \log x_2}{(1 - x_1)(1 - x_2)} - \frac{2 \log(x_1 x_2)}{(1 - x_1)(1 - x_1 x_2)} \right\} dx_1 dx_2. \end{aligned}$$

Since the integrand is equal to

$$-\frac{1 + x_1}{(1 - x_1)(1 - x_1 x_2)} \log x_1 + \frac{1 + x_2}{(1 - x_2)(1 - x_1 x_2)} \log x_2,$$

the right-hand side of the above equation is zero. This remark is related to the research for harmonic sums and Mellin transforms studied by the physicists Blümlein and Vermaseren [1, 18].

6 The number of the relations

For $m \geq 0$ we define V_m to be the $\mathbb{Q}$ -vector space whose basis is I_m , where $I_0 = \{\phi\}$. In this section, we give the dimension of the space $u\sigma(V * V) \cap V_m$, which is the number of independent relations among MZV's of weight m obtained from Corollary 5.5. Since $u\sigma$ is an isomorphism of the graded vector space $\tilde{V} = \bigoplus_{m \geq 0} V_m$, we have

$$\dim_{\mathbb{Q}}(u\sigma(V * V) \cap V_m) = \dim_{\mathbb{Q}}((V * V) \cap V_m). \quad (18)$$

Since the bilinear mapping $*$: $\tilde{V} \times \tilde{V} \rightarrow \tilde{V}$ is associative and commutative, $(\tilde{V}, *)$ is a commutative $\mathbb{Q}$ -algebra with unit ϕ . The key to proving Corollary 6.5 is the fact that $(\tilde{V}, *)$ is the polynomial algebra on the set of Lyndon words over $\mathbb{Z}_{\geq 1}$, which is due to Malvenuto and Reutenauer [12].

We start with the definition of the Lyndon words. Let A be a set. A word over A is a finite sequence of elements of A . We denote the length of a word u by $l(u)$, that is, we put $l(u) = p$ for a word $u = (a_1, \dots, a_p)$ ($a_i \in A$). For two words $u = (a_1, \dots, a_p)$ and $v = (b_1, \dots, b_q)$, we define a word uv by

$$uv = (a_1, \dots, a_p, b_1, \dots, b_q).$$

A word $(a_1, a_2, \dots, a_p)$ is usually written as $a_1 a_2 \dots a_p$. A lexicographic order on the set of words is given by a total order $\preceq$ on A : For any words $u = a_1 \dots a_p$, $v = b_1 \dots b_q$, one sets $u \preceq v$ if either

$$p \leq q \quad \text{and} \quad a_1 = b_1, \dots, a_p = b_p$$

or there exists some $1 \leq i \leq \min\{p, q\}$ such that

$$a_1 = b_1, \dots, a_{i-1} = b_{i-1}, a_i \prec b_i.$$

Definition 6.1. Let $(A, \preceq)$ be a totally ordered set. A word over A is called a Lyndon word if it is strictly smaller than any of its proper right factors, that is, $u = a_1 \dots a_p$ ($a_i \in A$) is a Lyndon word if $u \prec a_i \dots a_p$ for any $2 \leq i \leq p$.

We denote by $\psi_k(n)$ the number of Lyndon words of length n over a finite set of cardinality k . Then we have

$$k^n = \sum_{d|n} d\psi_k(d) \quad (19)$$

and

$$n\psi_k(n) = \sum_{d|n} \mu\left(\frac{n}{d}\right) k^d, \quad (20)$$

where the sums run over the positive divisors of n and μ is the Möbius function (see [11, p. 65]). The equality (20) is obtained from (19) and the Möbius inversion formula.

In Section 2, we defined a multi-index to be a finite sequence of positive integers. Namely, a multi-index is nothing else but a word over $\mathbb{Z}_{\geq 1}$. Let $\mathbb{Z}_{\geq 1}$

be totally ordered by the usual relation. We denote the set of Lyndon words over $\mathbb{Z}_{\geq 1}$ by L and the set of elements in L of weight m by L_m . We define $\Psi(m)$ to be the cardinality of the set L_m .

Proposition 6.2. *We have $\Psi(m) = \psi_2(m)$ for any $m \geq 2$.*

Proof. Let W_m be the set of words over $\{x, y\}$ of length m which start with x . The elements of I_m are in one-to-one correspondence with those of W_m by the mapping

$$(\mu_1, \mu_2, \dots, \mu_p) \mapsto x \underbrace{y \cdots y}_{\mu_1 - 1} x \underbrace{y \cdots y}_{\mu_2 - 1} \cdots x \underbrace{y \cdots y}_{\mu_p - 1}.$$

If we define a total order on the set $\{x, y\}$ by setting $x \prec y$, we can easily show that the Lyndon words in I_m are in one-to-one correspondence with the Lyndon words in W_m under the above correspondence. A Lyndon word over $\{x, y\}$ whose length is greater than or equal to 2 necessarily starts with x . Hence the assertion follows. $\square$

Corollary 6.3. *We have*

$$\Psi(m) = \frac{1}{m} \sum_{d|m} \mu\left(\frac{m}{d}\right) 2^d$$

for any $m \geq 2$.

Proof. It follows from (20) and Proposition 6.2. $\square$

For proofs of the following theorem, see [5, Section 2] or [12, Section 2].

Theorem 6.4. *$\tilde{V}$ is the polynomial algebra over $\mathbb{Q}$ on the set L .*

Corollary 6.5. *We have*

$$\dim_{\mathbb{Q}}(u\sigma(V * V) \cap V_m) = 2^{m-1} - \frac{1}{m} \sum_{d|m} \mu\left(\frac{m}{d}\right) 2^d$$

for any $m \geq 2$.

Proof. By Theorem 6.4, we have

$$V_m = \mathbb{Q}L_m \oplus ((V * V) \cap V_m)$$

for any $m \geq 1$. Since $\dim_{\mathbb{Q}} V_m = 2^{m-1}$, the assertion follows from Corollary 6.3 and (18). $\square$

7 The duality

In this section, we prove that the assertion of Corollary 5.5 contains the duality. For non-negative integers p, q with $(p, q) \neq (0, 0)$, we define $\mathcal{D}_{p,q}$ to be the set of $2 \times l$ ($l \geq 1$) matrices (k_{ij}) with entries in the non-negative integers which satisfy $k_{1j} \leq 1$ ($1 \leq j \leq l$), $k_{1j} + k_{2j} > 0$ ($1 \leq j \leq l$), $\sum_{j=1}^l k_{1j} = p$ and $\sum_{j=1}^l k_{2j} = q$. For example, $\mathcal{D}_{1,2}$ consists of

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 0 \end{pmatrix}, \\ \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 2 & 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix}. \quad (21)$$

We put

$$\mathcal{D}_{p,q}^0 = \{(k_{ij}) \in \mathcal{D}_{p,q} \mid k_{11} = 0\} \quad \text{and} \quad \mathcal{D}_{p,q}^1 = \{(k_{ij}) \in \mathcal{D}_{p,q} \mid k_{11} = 1\}.$$

For $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p)$, $\boldsymbol{\nu} = (\nu_1, \dots, \nu_q) \in \tilde{I}$ with $(\boldsymbol{\mu}, \boldsymbol{\nu}) \neq (\phi, \phi)$, $l(\boldsymbol{\mu}) = p$ and $l(\boldsymbol{\nu}) = q$, we define a mapping

$$\Phi_{p,q;\boldsymbol{\mu},\boldsymbol{\nu}}: \mathcal{D}_{p,q} \rightarrow \bigcup_{\boldsymbol{\beta} \leq \boldsymbol{\nu}} \mathcal{H}_{\boldsymbol{\mu},\boldsymbol{\beta}}, \quad \begin{pmatrix} k_{11} & \cdots & k_{1l} \\ k_{21} & \cdots & k_{2l} \end{pmatrix} \mapsto \begin{pmatrix} m_{11} & \cdots & m_{1l} \\ m_{21} & \cdots & m_{2l} \end{pmatrix}$$

by

$$\begin{aligned} & (m_{11}, m_{12}, \dots, m_{1l}) \\ &= (\underbrace{\mu_1 + \mu_2 + \cdots + \mu_x}_{k_{11}}, \underbrace{\mu_{x+1} + \mu_{x+2} + \cdots + \mu_y}_{k_{12}}, \dots, \underbrace{\mu_{z+1} + \mu_{z+2} + \cdots + \mu_p}_{k_{1l}}) \end{aligned}$$

and

$$\begin{aligned} & (m_{21}, m_{22}, \dots, m_{2l}) \\ &= (\underbrace{\nu_1 + \nu_2 + \cdots + \nu_{x'}}_{k_{21}}, \underbrace{\nu_{x'+1} + \nu_{x'+2} + \cdots + \nu_{y'}}_{k_{22}}, \dots, \underbrace{\nu_{z'+1} + \nu_{z'+2} + \cdots + \nu_q}_{k_{2l}}), \end{aligned}$$

where $\mathcal{H}_{\boldsymbol{\mu},\boldsymbol{\beta}}$ is defined in Section 2. For example, $\Phi_{1,2;\boldsymbol{\mu},\boldsymbol{\nu}}$ sends elements of (21) to

$$\begin{pmatrix} \mu_1 & 0 & 0 \\ 0 & \nu_1 & \nu_2 \end{pmatrix}, \begin{pmatrix} 0 & \mu_1 & 0 \\ \nu_1 & 0 & \nu_2 \end{pmatrix}, \begin{pmatrix} 0 & 0 & \mu_1 \\ \nu_1 & \nu_2 & 0 \end{pmatrix}, \\ \begin{pmatrix} \mu_1 & 0 \\ \nu_1 & \nu_2 \end{pmatrix}, \begin{pmatrix} 0 & \mu_1 \\ \nu_1 & \nu_2 \end{pmatrix}, \begin{pmatrix} \mu_1 & 0 \\ 0 & \nu_1 + \nu_2 \end{pmatrix}, \begin{pmatrix} 0 & \mu_1 \\ \nu_1 + \nu_2 & 0 \end{pmatrix}, \begin{pmatrix} \mu_1 \\ \nu_1 + \nu_2 \end{pmatrix},$$

respectively. It is easily seen that $\Phi_{p,q;\boldsymbol{\mu},\boldsymbol{\nu}}$ is a bijection.

Proposition 7.1. For any $\mu = (\mu_1, \dots, \mu_p) \in I$, we have

$$\sum_{h=0}^p (-1)^h \tau(\mu_1, \dots, \mu_h) * d(\mu_{h+1}, \dots, \mu_p) = 0,$$

where $(\mu_1, \dots, \mu_h) = \phi$ if $h = 0$ and $(\mu_{h+1}, \dots, \mu_p) = \phi$ if $h = p$.

Proof. Let $0 \leq h \leq p$. We denote the composition

$$\begin{aligned} \mathcal{D}_{h,p-h} &\xrightarrow{\Phi_{h,p-h;\tau(\mu_1, \dots, \mu_h), (\mu_{h+1}, \dots, \mu_p)}} \bigcup_{\beta \leq (\mu_{h+1}, \dots, \mu_p)} \mathcal{H}_{\tau(\mu_1, \dots, \mu_h), \beta} \rightarrow I \\ \begin{pmatrix} k_{11} & \cdots & k_{1l} \\ k_{21} & \cdots & k_{2l} \end{pmatrix} &\mapsto \begin{pmatrix} m_{11} & \cdots & m_{1l} \\ m_{21} & \cdots & m_{2l} \end{pmatrix} \mapsto (m_{11} + m_{21}, \dots, m_{1l} + m_{2l}) \end{aligned}$$

by ψ_h . By the definition of the harmonic product $*$ and the mapping d , we have

$$\tau(\mu_1, \dots, \mu_h) * d(\mu_{h+1}, \dots, \mu_p) = \sum_{D \in \mathcal{D}_{h,p-h}} \psi_h(D).$$

For any $0 \leq h < p$, the mapping

$$\begin{aligned} \mathcal{D}_{h,p-h}^0 &\rightarrow \mathcal{D}_{h+1,p-h-1}^1, \\ D = \begin{pmatrix} 0 & k_{12} & \cdots & k_{1l} \\ k_{21} & k_{22} & \cdots & k_{2l} \end{pmatrix} &\mapsto D' = \begin{pmatrix} 1 & k_{12} & \cdots & k_{1l} \\ k_{21} - 1 & k_{22} & \cdots & k_{2l} \end{pmatrix} \end{aligned}$$

is a bijection and clearly $\psi_h(D) = \psi_{h+1}(D')$. Consequently, we have

$$\begin{aligned} &\sum_{h=0}^p (-1)^h \tau(\mu_1, \dots, \mu_h) * d(\mu_{h+1}, \dots, \mu_p) \\ &= \sum_{h=0}^p (-1)^h \left(\sum_{D \in \mathcal{D}_{h,p-h}^0} + \sum_{D \in \mathcal{D}_{h,p-h}^1} \right) \psi_h(D) \\ &= \sum_{h=0}^{p-1} (-1)^h \sum_{D \in \mathcal{D}_{h,p-h}^0} \psi_h(D) + \sum_{h=0}^{p-1} (-1)^{h+1} \sum_{D \in \mathcal{D}_{h+1,p-h-1}^1} \psi_{h+1}(D) \\ &= 0. \end{aligned}$$

Thus we get the proposition. $\square$

The duality reads

$$(\tau - *)(V) \subset \ker \zeta^+.$$

Therefore Corollary 7.2 says that the assertion of Corollary 5.5 contains the duality.

Corollary 7.2. We have $(\tau - *)(V) \subset u\sigma(V * V)$.

Proof. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$. By Proposition 7.1, we have

$$-\sum_{h=1}^{p-1} (-1)^h \tau(\mu_1, \dots, \mu_h) * d(\mu_{h+1}, \dots, \mu_p) = (\sigma\tau + d)(\boldsymbol{\mu}). \quad (22)$$

By Proposition 2.1 and 2.3 (v), we have $u\sigma(\sigma\tau + d) = (\tau - *)u$. Applying $u\sigma$ to both sides of (22), we obtain $(\tau - *)u(\boldsymbol{\mu}) \in u\sigma(V * V)$. Thus we have completed the proof. $\square$

8 Ohno's relation

In this section, we prove that the assertion of Corollary 5.5 contains Ohno's relation. In Section 2, we defined $\tilde{V}$ to be the $\mathbb{Q}$ -vector space whose basis is $\tilde{I} := \{\phi\} \cup I$. For $v \in \tilde{V}$, we define the $\mathbb{Q}$ -linear mapping m_v from $\tilde{V}$ to $\tilde{V}$ by

$$m_v(w) = v * w \quad (w \in \tilde{V}).$$

For $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$, we define the mapping $O_{\boldsymbol{\mu}}: \tilde{I} \rightarrow \tilde{V}$ by

$$O_{\boldsymbol{\mu}}(\boldsymbol{\nu}) = \sum_{1 \leq i_1 < \dots < i_p \leq q} (\nu_1, \dots, \nu_{i_1} + \mu_1, \dots, \nu_{i_p} + \mu_p, \dots, \nu_q)$$

($\boldsymbol{\nu} = (\nu_1, \dots, \nu_q) \in I$) and $O_{\boldsymbol{\mu}}(\phi) = 0$. We extend the mapping $O_{\boldsymbol{\mu}}$ linearly on $\tilde{V}$. By setting $O_{\phi} = \text{id}_{\tilde{V}}$, we obtain the mapping $\tilde{I} \rightarrow \text{End}_{\mathbb{Q}}(\tilde{V})$, $\boldsymbol{\mu} \mapsto O_{\boldsymbol{\mu}}$. We extend this mapping linearly on $\tilde{V}$. We define $\overline{O}_v = *O_v*$ for any $v \in \tilde{V}$.

Proposition 8.1. *For any $\boldsymbol{\mu} \in \tilde{I}$, we have $O_{\boldsymbol{\mu}}\sigma = \sigma O_{\boldsymbol{\mu}}$ and $\overline{O}_{\boldsymbol{\mu}}\sigma = (-1)^{|\boldsymbol{\mu}|}\sigma\overline{O}_{\boldsymbol{\mu}}$.*

Proof. The first assertion is easily seen. Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_p) \in I$. Let $\boldsymbol{\nu} \in I$ and $\boldsymbol{\nu}^* = (\nu_1^*, \dots, \nu_r^*)$. Then by (3) we have

$$\sigma\overline{O}_{\boldsymbol{\mu}}(\boldsymbol{\nu}) = \sum_{1 \leq i_1 < \dots < i_p \leq r} (-1)^{|\boldsymbol{\mu}| + |\boldsymbol{\nu}| + 1 - r} * (\nu_1^*, \dots, \nu_{i_1}^* + \mu_1, \dots, \nu_{i_p}^* + \mu_p, \dots, \nu_r^*).$$

Since $|\boldsymbol{\nu}| + 1 - r = l(\boldsymbol{\nu})$, the second assertion follows. $\square$

Proposition 8.2. *For any $v \in \tilde{V}$, we have $O_v\tau = \tau O_{\tau(v)}$.*

Proof. This is easily seen. $\square$

Proposition 8.3. *For any $v, w \in \tilde{V}$, we have $O_v O_w = O_{v * w}$.*

Proof. It is easily seen that $O_{\boldsymbol{\mu}} O_{\boldsymbol{\nu}} = O_{\boldsymbol{\mu} * \boldsymbol{\nu}}$ for any $\boldsymbol{\mu}, \boldsymbol{\nu} \in \tilde{I}$. $\square$

In the following, if $r = 0$, we put

$$(r) = \phi \quad \text{and} \quad \underbrace{(1, \dots, 1)}_r = \phi.$$

Proposition 8.4. For any $\mu, \nu \in I$ and any integer $r \geq 0$, we have

$$O_{u(r)}(\mu \# \nu) = \sum_{\substack{k+l=r \\ k,l \geq 0}} O_{u(k)}(\mu) \# O_{u(l)}(\nu)$$

and

$$\overline{O}_{u(r)}(\mu \dot{\#} \nu) = \sum_{\substack{k+l=r \\ k,l \geq 0}} \overline{O}_{u(k)}(\mu) \dot{\#} \overline{O}_{u(l)}(\nu).$$

Proof. The first is easily seen. The second follows from the first and Proposition 2.6. $\square$

Proposition 8.5. Let $\mu \in I$. For any $\nu \in I$, we have

$$O_{\mu}((1) \dot{\#} \nu) = (1) \dot{\#} O_{\mu}(\nu) \quad \text{and} \quad \overline{O}_{\mu}((1) \# \nu) = (1) \# \overline{O}_{\mu}(\nu).$$

Proof. The first is easily seen. The second follows from the first and Proposition 2.6. $\square$

Proposition 8.6. For any $\mu \in I$ and any integer $r \geq 0$, we have

$$O_{\underbrace{1, \dots, 1}_r}(\mu) = \sum_{\substack{\nu^1 \# \dots \# \nu^{r+1} = \mu \\ \nu^1, \dots, \nu^r \in I, \nu^{r+1} \in \tilde{I}}} \nu^1 \dot{\#} ((1) \# \nu^2) \dot{\#} \dots \dot{\#} ((1) \# \nu^{r+1})$$

and

$$O_{u(r)}(\mu) = \sum_{\substack{\nu^1 \# \dots \# \nu^{r+1} = \mu \\ \nu^1, \dots, \nu^r \in \tilde{I}, \nu^{r+1} \in I}} (\nu^1 \# (1)) \dot{\#} \dots \dot{\#} (\nu^r \# (1)) \dot{\#} \nu^{r+1}.$$

Proof. These are easily seen. $\square$

By Propositions 8.6 and 2.6, we obtain

$$\overline{O}_{u(r)}(\mu) = \sum_{\substack{\nu^1 \dot{\#} \dots \dot{\#} \nu^{r+1} = \mu \\ \nu^1, \dots, \nu^r \in \tilde{I}, \nu^{r+1} \in I}} (\nu^1)^+ \# \dots \# (\nu^r)^+ \# \nu^{r+1}. \quad (23)$$

Ohno's relation reads

$$O_{u(r)}(t - *) (V) \subset \ker \zeta^+ \quad (r \geq 0).$$

The key to proving Corollary 8.12, which says that the assertion of Corollary 5.5 contains Ohno's relation, is the following proposition.

Proposition 8.7. For any integer $r \geq 0$, we have

$$u^{-1} m_{\underbrace{1, \dots, 1}_r} u = \sum_{k=0}^r \overline{O}_{u(r-k)} O_{\underbrace{1, \dots, 1}_k}.$$

Proof. It is immediate from Lemmas 8.8 and 8.10, which are proved later. $\square$

We associate a multi-index $\mu = (\mu_1, \dots, \mu_p)$ with a diagram

$$\begin{array}{ccccccc} \circ & \circ & \cdots & \circ & \downarrow & \circ & \circ & \cdots & \circ & \downarrow & \cdots & \downarrow & \circ & \circ & \cdots & \circ \\ \hline & & & \mu_1 & & & \mu_2 & & & \cdots & & & \mu_p & & & \end{array} \quad \left. \vphantom{\begin{array}{c} \circ \\ \downarrow \end{array}} \right\} \text{arrows},$$

where the number of arrows is $p - 1$. For example, we associate multi-indices $(2, 2)$ and $(1, 2, 1)$ with diagrams

$$\begin{array}{ccccccc} & & \downarrow & & & \downarrow & & \downarrow \\ \circ & \circ & \circ & \circ & \text{and} & \circ & \circ & \circ & \circ \end{array}.$$

Then multi-indices are in one-to-one correspondence with these diagrams. Let $n_1, \dots, n_r$ ($r \geq 1$) be non-negative integers with $n_1 + \dots + n_r = |\mu|$. We define $\nu^1, \dots, \nu^r \in \tilde{I}$ by

$$\begin{array}{ccccccc} \overbrace{}^{\nu^1} & \overbrace{}^{\nu^2} & & \overbrace{}^{\nu^r} & & & \\ \circ \cdots \circ & \circ \cdots \circ & \cdots & \circ \cdots \circ & & & \\ \hline & n_1 & & n_2 & & \cdots & n_r \end{array} \quad \left. \vphantom{\begin{array}{c} \circ \\ \downarrow \end{array}} \right\} \text{arrows of } \mu,$$

where the weight of ν^i is n_i . We call $(\nu^1, \dots, \nu^r)$ the $(n_1, \dots, n_r)$ -partition of μ and write $(\nu^1, \dots, \nu^r) \parallel \mu$. For example, the $(2, 1, 3)$ -partition and the $(3, 0, 3)$ -partition of $(2, 2, 2)$ are $((2), (1), (1, 2))$ and $((2, 1), \phi, (1, 2))$ by the following diagrams

$$\begin{array}{ccccccc} & & \downarrow & & \downarrow & & \downarrow \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \\ \hline & 2 & & 1 & & 3 & \end{array} \quad \text{and} \quad \begin{array}{ccccccc} & & \downarrow & & \downarrow & & \downarrow \\ \circ & \circ & \circ & \circ & \circ & \circ & \circ \\ \hline & 3 & & & & 3 & \end{array}.$$

Lemma 8.8. For any $\mu \in I$ and any integer $r \geq 0$, we have

$$u^{-1} m_{\underbrace{1, \dots, 1}_r} u(\mu) = \sum_{\substack{(\nu^1, \dots, \nu^{r+1}) \parallel \mu \\ \nu^i \in \tilde{I}}} (\nu^1)^+ \# \cdots \# (\nu^r)^+ \# \nu^{r+1}.$$

Proof. We define a mapping $f: \tilde{I} \rightarrow V$ by $f(\phi) = (1)$ and $f(\alpha) = \alpha \# (1) + \alpha \# (1)$ ($\alpha \in I$). Then we have

$$m_{\underbrace{1, \dots, 1}_r}(\mu) = \sum_{\substack{\alpha^1 \# \cdots \# \alpha^{r+1} = \mu \\ \alpha^i \in \tilde{I}}} f(\alpha^1) \# \cdots \# f(\alpha^r) \# \alpha^{r+1}$$

and therefore

$$m_{\underbrace{1, \dots, 1}_r} u(\mu) = \sum_{\substack{\alpha^1 \# \cdots \# \alpha^{r+1} \geq \mu \\ \alpha^i \in \tilde{I}}} f(\alpha^1) \# \cdots \# f(\alpha^r) \# \alpha^{r+1}. \quad (24)$$

We construct a mapping

$$\begin{aligned} & \{(\alpha^1, \dots, \alpha^{r+1}) \in \tilde{I}^{r+1} \mid \alpha^1 \# \dots \# \alpha^{r+1} \geq \mu\} \rightarrow \\ & \left\{ ((\alpha^1, \dots, \alpha^{r+1}), (\nu^1, \dots, \nu^{r+1})) \in \tilde{I}^{r+1} \times \tilde{I}^{r+1} \mid \begin{array}{l} (\nu^1, \dots, \nu^{r+1}) \parallel \mu, \\ \alpha^i \geq \nu^i \ (1 \leq i \leq r+1) \end{array} \right\}, \\ & (\alpha^1, \dots, \alpha^{r+1}) \mapsto ((\alpha^1, \dots, \alpha^{r+1}), (\nu^1, \dots, \nu^{r+1})) \end{aligned}$$

by defining $(\nu^1, \dots, \nu^{r+1})$ to be the $(|\alpha^1|, \dots, |\alpha^{r+1}|)$ -partition of μ . The mapping

$$((\alpha^1, \dots, \alpha^{r+1}), (\nu^1, \dots, \nu^{r+1})) \mapsto (\alpha^1, \dots, \alpha^{r+1})$$

is its inverse and thereby this mapping is a bijection. Hence, the right-hand side of (24) is equal to

$$\begin{aligned} & \sum_{\substack{(\nu^1, \dots, \nu^{r+1}) \parallel \mu \\ \nu^i \in \tilde{I}}} \sum_{\substack{\alpha^1 \geq \nu^1 \\ \dots \\ \alpha^{r+1} \geq \nu^{r+1}}} f(\alpha^1) \# \dots \# f(\alpha^r) \# \alpha^{r+1} \\ &= \sum_{\substack{(\nu^1, \dots, \nu^{r+1}) \parallel \mu \\ \nu^i \in \tilde{I}}} \left\{ \sum_{\alpha^1 \geq \nu^1} f(\alpha^1) \right\} \# \dots \# \left\{ \sum_{\alpha^r \geq \nu^r} f(\alpha^r) \right\} \# \left\{ \sum_{\alpha^{r+1} \geq \nu^{r+1}} \alpha^{r+1} \right\} \\ &= \sum_{\substack{(\nu^1, \dots, \nu^{r+1}) \parallel \mu \\ \nu^i \in \tilde{I}}} u((\nu^1)^+) \# \dots \# u((\nu^r)^+) \# u(\nu^{r+1}). \end{aligned}$$

By Proposition 2.7, we obtain the assertion. $\square$

Remark 8.9. We have $u^{-1} = \sigma u \sigma$ by Proposition 2.3 (iii). According to Corollary 5.5, we obtain

$$\sigma \sum_{\substack{(\nu^1, \dots, \nu^{r+1}) \parallel \mu \\ \nu^i \in \tilde{I}}} (\nu^1)^+ \# \dots \# (\nu^r)^+ \# \nu^{r+1} \in \ker \zeta^+$$

for any $\mu \in I$ and any integer $r \geq 1$. For example, for $\mu = (2, 1)$ and $r = 2$, all of partitions (ν^1, ν^2) of μ are $((2), (1))$, $((1), (1, 1))$, $((2, 1), \phi)$, $(\phi, (2, 1))$ by the diagrams

$$\begin{array}{ccc} \downarrow & & \downarrow \\ \bigcirc & \bigcirc & \bigcirc \\ \hline \end{array}, \quad \begin{array}{ccc} \downarrow & & \downarrow \\ \bigcirc & \bigcirc & \bigcirc \\ \hline \end{array} \quad \text{and} \quad \begin{array}{ccc} \downarrow & & \downarrow \\ \bigcirc & \bigcirc & \bigcirc \\ \hline \end{array}.$$

Since $(2)^+ \# (1) = (3, 1)$, $(1)^+ \# (1, 1) = (2, 1, 1)$, $(2, 1)^+ \# \phi = (2, 2)$ and $\phi^+ \# (2, 1) = (1, 2, 1)$, we obtain $(3, 1) - (2, 1, 1) + (2, 2) - (1, 2, 1) \in \ker \zeta^+$.

Lemma 8.10. *For any $\mu \in I$ and any integer $r \geq 0$, we have*

$$\sum_{k=0}^r \overline{O}_{u(r-k)} \underbrace{O_{1, \dots, 1}}_k(\mu) = \sum_{\substack{(\nu^1, \dots, \nu^{r+1}) \parallel \mu \\ \nu^i \in \tilde{I}}} (\nu^1)^+ \# \dots \# (\nu^r)^+ \# \nu^{r+1}.$$

Proof. By Proposition 8.6, we have

$$O_{\underbrace{1, \dots, 1}_{k-1}}(\mu) = \sum_{\substack{\alpha^1 \# \dots \# \alpha^k = \mu \\ \alpha^1, \dots, \alpha^{k-1} \in I, \alpha^k \in \tilde{I}}} \alpha^1 \dot{\#} ((1) \# \alpha^2) \dot{\#} \dots \dot{\#} ((1) \# \alpha^k).$$

Therefore, by Propositions 8.4 and 8.5, we have

$$\begin{aligned} \overline{O}_{u(r-k+1)} O_{\underbrace{1, \dots, 1}_{k-1}}(\mu) &= \sum_{\substack{\alpha^1 \# \dots \# \alpha^k = \mu \\ \alpha^1, \dots, \alpha^{k-1} \in I, \alpha^k \in \tilde{I}}} \sum_{\substack{m_1 + \dots + m_k = r+1 \\ m_i \geq 1}} \\ &\quad \overline{O}_{u(m_1-1)}(\alpha^1) \dot{\#} ((1) \# \overline{O}_{u(m_2-1)}(\alpha^2)) \dot{\#} \dots \dot{\#} ((1) \# \overline{O}_{u(m_k-1)}(\alpha^k)), \end{aligned}$$

where we put

$$\overline{O}_{u(m_k-1)}(\alpha^k) = \underbrace{(1, \dots, 1)}_{m_k-1}$$

if $\alpha^k = \phi$. By (23), this is equal to

$$\begin{aligned} &\sum_{\substack{\alpha^1 \# \dots \# \alpha^k = \mu \\ \alpha^1, \dots, \alpha^{k-1} \in I, \alpha^k \in \tilde{I}}} \sum_{\substack{m_1 + \dots + m_k = r+1 \\ m_i \geq 1}} \\ &\quad \left\{ \sum_{\substack{\lambda^{11} \dot{\#} \dots \dot{\#} \lambda^{1m_1} = \alpha^1 \\ \lambda^{11}, \dots, \lambda^{1m_1-1} \in \tilde{I}, \lambda^{1m_1} \in I}} (\lambda^{11})^+ \# \dots \# (\lambda^{1m_1-1})^+ \# \lambda^{1m_1} \right\} \\ &\quad \dot{\#} \left\{ (1) \# \sum_{\substack{\lambda^{21} \dot{\#} \dots \dot{\#} \lambda^{2m_2} = \alpha^2 \\ \lambda^{21}, \dots, \lambda^{2m_2-1} \in \tilde{I}, \lambda^{2m_2} \in I}} (\lambda^{21})^+ \# \dots \# (\lambda^{2m_2-1})^+ \# \lambda^{2m_2} \right\} \\ &\quad \dot{\#} \dots \\ &\quad \dot{\#} \left\{ (1) \# \sum_{\substack{\lambda^{k1} \dot{\#} \dots \dot{\#} \lambda^{km_k} = \alpha^k \\ \lambda^{k1}, \dots, \lambda^{km_k} \in \tilde{I} \\ \alpha^k \neq \phi \Rightarrow \lambda^{km_k} \neq \phi}} (\lambda^{k1})^+ \# \dots \# (\lambda^{km_k-1})^+ \# \lambda^{km_k} \right\} \\ &= \sum_{\substack{\alpha^1 \# \dots \# \alpha^k = \mu \\ \alpha^1, \dots, \alpha^{k-1} \in I \\ \alpha^k \in \tilde{I}}} \sum_{\substack{m_1 + \dots + m_k = r+1 \\ m_i \geq 1}} \sum_{\substack{\lambda^{11} \dot{\#} \dots \dot{\#} \lambda^{1m_1} = \alpha^1 \\ \lambda^{11}, \dots, \lambda^{1m_1-1} \in \tilde{I} \\ \lambda^{1m_1} \in I}} \dots \sum_{\substack{\lambda^{k1} \dot{\#} \dots \dot{\#} \lambda^{km_k} = \alpha^k \\ \lambda^{k1}, \dots, \lambda^{km_k} \in \tilde{I} \\ \alpha^k \neq \phi \Rightarrow \lambda^{km_k} \neq \phi}} \\ &\quad (\lambda^{11})^+ \# \dots \# (\lambda^{1m_1})^+ \# \dots \# (\lambda^{k1})^+ \# \dots \# (\lambda^{km_k-1})^+ \# \lambda^{km_k}. \end{aligned}$$

We define a mapping

$$\left\{ (\nu^1, \dots, \nu^{r+1}) \in \tilde{I}^{r+1} \mid (\nu^1, \dots, \nu^{r+1}) \parallel \mu \right\} \rightarrow \left\{ ((\lambda^{11}, \dots, \lambda^{1m_1}), \dots, (\lambda^{k1}, \dots, \lambda^{km_k})) \mid \begin{array}{l} 1 \leq k \leq r+1, m_1 + \dots + m_k = r+1, m_i \geq 1, \\ (\lambda^{11} \# \dots \# \lambda^{1m_1}) \# \dots \# (\lambda^{k1} \# \dots \# \lambda^{km_k}) = \mu, \\ \lambda^{ij} \in \tilde{I}, \lambda^{1m_1}, \dots, \lambda^{k-1m_{k-1}} \neq \phi, \lambda^{km_k} = \phi \Rightarrow \lambda^{k1} = \dots = \lambda^{km_k} = \phi \end{array} \right\},$$

$$(\nu^1, \dots, \nu^{r+1}) \mapsto ((\lambda^{11}, \dots, \lambda^{1m_1}), \dots, (\lambda^{k1}, \dots, \lambda^{km_k}))$$

by


$$(\lambda^{11}, \dots, \lambda^{1m_1}, \dots, \lambda^{k1}, \dots, \lambda^{km_k}) = (\nu^1, \dots, \nu^{r+1})$$

and

$$\{m_1, m_1 + m_2, \dots, m_1 + m_2 + \dots + m_k\}$$

$$= \left\{ 1 \leq i \leq r+1 \mid \begin{array}{l} |\nu^1| + \dots + |\nu^i| = \mu_1 + \dots + \mu_j \\ \text{for some } 1 \leq j \leq l(\mu) \text{ and } \nu^i \neq \phi \text{ if } i \neq r+1 \end{array} \right\}.$$

The following diagram illustrates the definition of this mapping.



($M_i = m_1 + \dots + m_i$. There exist no arrows of μ at the positions of $\times$.)

The inverse is given by

$$((\lambda^{11}, \dots, \lambda^{1m_1}), \dots, (\lambda^{k1}, \dots, \lambda^{km_k}))$$

$$\mapsto (\lambda^{11} \# \dots \# \lambda^{1m_1}) \# \dots \# (\lambda^{k1} \# \dots \# \lambda^{km_k}),$$

and therefore the mapping is a bijection. As a result, we obtain

$$\sum_{k=1}^{r+1} \bar{O}_{u(r-k+1)} \underbrace{O_{1, \dots, 1}}_{k-1}(\mu) = \sum_{\substack{(\nu^1, \dots, \nu^{r+1}) \parallel \mu \\ \nu^i \in \tilde{I}}} (\nu^1)^+ \# \dots \# (\nu^r)^+ \# \nu^{r+1}.$$

We have thus completed the proof. $\square$

Proposition 8.11. *For any integer $r \geq 0$, we have*

$$\sum_{k=0}^r (-1)^k u^{-1} \underbrace{m_1, \dots, 1}_{r-k} u O_{u(k)} = \overline{O}_{u(r)}.$$

Proof. By Proposition 8.7, the left-hand side is equal to

$$\sum_{r \geq l \geq k \geq 0} (-1)^k \overline{O}_{u(r-l)} \underbrace{O_1, \dots, 1}_{l-k} O_{u(k)}. \quad (25)$$

By Propositions 8.3 and 7.1, if $l > 0$, we have

$$\sum_{k=0}^l (-1)^k \underbrace{O_1, \dots, 1}_{l-k} O_{u(k)} = \sum_{k=0}^l (-1)^k O(\underbrace{1, \dots, 1}_{l-k} * \underbrace{1, \dots, 1}_k) = 0.$$

Therefore (25) is equal to $\overline{O}_{u(r)}$. $\square$

Since $u^{-1} = \sigma u \sigma$, by Proposition 8.11 we have

$$\sigma(\overline{O}_{u(r)} - (-1)^r O_{u(r)}) = \sum_{k=0}^{r-1} (-1)^k u \sigma \underbrace{m_1, \dots, 1}_{r-k} u O_{u(k)}$$

and therefore

$$\sigma(\overline{O}_{u(r)} - (-1)^r O_{u(r)})(V) \subset u\sigma(V * V). \quad (26)$$

Corollary 8.12. *For any integer $r \geq 0$, we have $O_{u(r)}(\tau - *) (V) \subset u\sigma(V * V)$.*

Proof. Since

$$\sigma((-1)^r \overline{O}_{u(r)} - O_{u(r)})(V) = (\overline{O}_{u(r)} - O_{u(r)})(V)$$

by Proposition 8.1, we have

$$(\overline{O}_{u(r)} - O_{u(r)})(V) \subset u\sigma(V * V) \quad (27)$$

by (26). Since $O_{u(r)}\tau = \tau O_{u(r)}$ by Proposition 8.2, we have

$$O_{u(r)}\tau - O_{u(r)}* = (\tau - *)O_{u(r)} + (\overline{O}_{u(r)} - O_{u(r)})*.$$

This equality together with (27) and Corollary 7.2 implies the assertion of Corollary 8.12. $\square$

Acknowledgments. The author would like to express his gratitude to Professor Masanobu Kaneko and Professor Yasuo Ohno for valuable comments and suggestions. Especially Professor Ohno recommended the author to calculate the numerical values of the dimensions of our relations and Professor Kaneko suggested the dimension formula as is stated in the introduction. The author would also like to thank Professor Yoshio Tanigawa for helpful advices. Finally the author would like to thank the anonymous referee for a number of valuable comments to improve the presentation of the paper.

References

- [1] J. Blümlein and S. Kurth, Harmonic sums and Mellin transforms up to two-loop order, *Phys. Rev. D* **60** (1999), 014018.
- [2] L. Euler, *Meditationes circa singulare serierum genus*, *Novi Comm. Acad. Sci. Petropol.* **20** (1775), 140-186; reprinted in his *Opera Omnia* ser. I, vol. **15**, B. G. Teubner, Berlin (1927), 217-267.
- [3] A. O. Gel'fond, *Calculus of Finite Differences*, Hindustan, 1971.
- [4] M. Hoffman, Multiple harmonic series, *Pacific J. Math.*, **152** (1992), 275-290.
- [5] M. Hoffman, The algebra of multiple harmonic series, *J. Algebra* **194** (1997), 477-495.
- [6] M. Hoffman, Quasi-symmetric functions and mod p multiple harmonic sums, preprint arXiv:math.NT/0401319.
- [7] M. Hoffman and Y. Ohno, Relation of multiple zeta values and their algebraic expression, *J. Algebra* **262** (2003), 332-347.
- [8] K. Ihara, M. Kaneko and D. Zagier, Derivation relations and regularized double shuffle relations of multiple zeta values, *Compositio Math.* **142** (2006), 307-338.
- [9] S. Izumi, *Teisahô*, Iwanami, 1934 (in Japanese).
- [10] M. Kaneko, On an extension of the derivation relation for multiple zeta values, *The Conference on L -Functions*, edited by L. Weng and M. Kaneko, World Scientific (2007), 89-94.
- [11] M. Lothaire, *Combinatorics on Words*, Addison-Wesley, 1983.
- [12] C. Malvenuto and C. Reutenauer, Duality between Quasi-Symmetric Functions and the Solomon Descent Algebra, *J. Algebra* **177** (1995), 967-982.
- [13] L. M. Milne-Thomson, *The Calculus of Finite Differences*, Macmillan, 1933.
- [14] S. Muneta, Algebraic setup of non-strict multiple zeta values, preprint arXiv:0711.0252
- [15] Y. Ohno, A generalization of the duality and sum formulas on the multiple zeta values, *J. Number Theory* **74** (1999), 39-43.
- [16] Y. Ohno and N. Wakabayashi, Cyclic sum of multiple zeta values, *Acta Arith.* **123** (2006), 289-295.
- [17] T. Tanaka, On extended derivation relations for multiple zeta values, preprint arXiv:0710.4920

- [18] J. A. M. Vermaseren, Harmonic sums, Mellin transforms and integrals, Int. J. Mod. Phys. A **14** (1999), 2037-2076.
- [19] D. Zagier, Values of zeta functions and their applications, in First European Congress of Mathematics, Vol. II, Progr. Math. **120** (1994), 497-512.